


2016

Organic Production and IPM Guide for Strawberries



NYS IPM Publication No. 226



Cornell University
Cooperative Extension



Integrated Pest Management



New York State
Department of
Agriculture & Markets

2016 PRODUCTION GUIDE FOR ORGANIC STRAWBERRIES

Coordinating Editors

Juliet Carroll* (Cornell University, New York State IPM Program)
Marvin Pritts* (Cornell University, Section of Horticulture)
Catherine Heidenreich* (Cornell University, Section of Horticulture)

Contributors and Resources

Kerik Cox* (Cornell University, Section of Plant Pathology and Plant Microbe-Biology)
Greg Loeb* (Cornell University, Department of Entomology)
Michael Helms* (Cornell University, Pesticide Management Education Program)
Andrew Landers (Cornell University, Department of Entomology)
Paul Curtis (Cornell University, Department of Natural Resources)
Courtney Weber (Cornell University, Section of Horticulture)
Laura McDermott (Cornell Cooperative Extension, Eastern New York Commercial Horticulture Program)
Elizabeth Bihn (Cornell University, Department of Food Science)

**Pesticide Information and Regulatory Compliance*

Staff Writers

Mary Kirkwyland and Elizabeth Graeper Thomas (Cornell University, New York State IPM Program)
Theodora Buden (Cornell University, New York State IPM Program)

Special Appreciation

Format based on the Cornell University Pest Management Guidelines for Berry Crops <https://ipmguidelines.org/>, content editor Marvin Pritts and coordinating editor Cathy Heidenreich; and on the Production Guide for Organic Grapes, coordinating editors Tim Weigle and Juliet Carroll.

Dedication

This publication is dedicated to Cathy Heidenreich who was tragically killed in an automobile accident in December 2014. Cathy put her heart and soul into her work, which included many hours on this guide.

Funded in part by the New York State Department of Agriculture and Markets.

The guidelines in this bulletin reflect the current authors' best effort to interpret a complex body of scientific research, and to translate this into practical management options. Following the guidance provided in this bulletin does not assure compliance with any applicable law, rule, regulation or standard, or the achievement of particular discharge levels from agricultural land.

Every effort has been made to provide correct, complete, and up-to-date pest management information for New York State at the time this publication was released for printing (June 2016). Changes in pesticide registrations, regulations, and guidelines occurring after publication are available in county Cornell Cooperative Extension offices or from the Pesticide Management Education Program web site (pmep.cce.cornell.edu). Trade names used herein are for convenience only. No endorsement of products is intended, nor is criticism of unnamed products implied.

This guide is not a substitute for pesticide labeling. Always read the product label before applying any pesticide.

Updates and additional information for this guide are available at www.nysipm.cornell.edu/organic_guide/. Please submit comments or suggested changes for these guides to organicguides@gmail.com.

How to cite this publication: Carroll, J., Pritts, M.P., and Heidenreich, C., eds. (2016). Production Guide for Organic Strawberries. New York State Integrated Pest Management Program. Ithaca, NY. 64 pages.
--

This guide is published by the New York State Integrated Pest Management Program, which is funded through Cornell University, Cornell Cooperative Extension, the New York State Department of Agriculture and Markets, the New York State Department of Environmental Conservation, and USDA-NIFA. Cornell Cooperative Extension provides equal program and employment opportunities. NYS IPM Publication number 226, June 2016. http://www.nysipm.cornell.edu/organic_guide/ .

Table of Contents

1. GENERAL ORGANIC MANAGEMENT PRACTICES	3
2. SOIL HEALTH	4
3. SITE SELECTION	5
3.1 Organic Certification Site Requirements.....	5
3.2 Soil and Air Drainage and Soil Depth	5
3.3 Soil Testing.....	6
3.4 Previous Cropping History.....	6
3.5 Irrigation Water Source	6
4. COVER CROPS	7
4.1 Goals and Timing for Cover Crops	7
4.2 Legumes.....	9
5. VARIETY SELECTION	9
6. NUTRIENT MANAGEMENT.....	10
6.1 Soil and Leaf Analysis.....	11
6.2 Soil pH	12
6.3 Managing Nutrients.....	12
6.4 Preparing a Nitrogen Budget.....	13
7. ORGANIC STRAWBERRY IPM.....	15
7.1 Developing a Strawberry IPM Strategy	15
7.2 Weed Management.....	16
7.3 Principles of Insect and Disease Management	17
7.4 DISEASES OF PRIMARY CONCERN	19
7.5 OTHER DISEASES OF NOTE	34
7.6 INSECTS AND MITES OF PRIMARY CONCERN	36
7.7 MINOR AND SPORADIC INSECT AND MITE PESTS	45
7.8 Slug Management (<i>various species</i>)	56
7.9 Wildlife Management.....	57
7.10 Considerations During Harvest and Renovation.....	58
8. FOOD SAFETY	59
9. SMALL-SCALE SPRAYER TECHNOLOGY	60
9.1 Spraying Small Strawberry Plantings	60
9.2 Selecting a Small Sprayer for the Small, Organic Strawberry Planting	62
10. PESTICIDES MENTIONED IN THIS PUBLICATION	63
10.1 Pesticides Registered for use in Organic Strawberry Production.....	66
10.2 Pesticide Regulatory Considerations	66
10.3 Optimizing Pesticide Effectiveness	67
11. REFERENCES AND RESOURCES	67
11. GLOSSARY.....	69

INTRODUCTION

This guide for organic strawberry production is focused on nutrient and pest management practices and includes topics that have an impact on improving plant health and reducing pest problems. The guide is divided into sections, but the interrelated quality of organic cropping systems makes each section relevant to the others.

Strawberries are moderately amenable to organic production. The greatest challenge, by far, is weeds, particularly in the planting year. Studies have shown that sustained weed pressure in the planting year can negatively affect yield for several subsequent years. It is also difficult to provide a large amount of nitrate nitrogen when the strawberry plant needs it most: early spring and late fall. There are also a few pests that can be difficult to control organically if the weather does not cooperate (e.g. gray mold and anthracnose fruit rots). Use of protected production structures, such as low tunnels, may be needed to adequately manage these diseases. That said, with sufficient attention to weed management, especially in the planting year, and with good soil nitrogen reserves, strawberries can be successfully grown with organic production methods.

Organic strawberry production systems generally share five common characteristics, described in the Strawberry Production Guide for the Northeast, Midwest, and Eastern Canada, NRAES-88:

1. Several years elapse between successive strawberry crops. That is, practice 3- to 5-year-long crop rotations.
2. The production cycle is short, only one or two fruiting years, to avoid the establishment of perennial weeds and depletion of nitrogen reserves.
3. The labor requirements are high because of the need for hand-weeding and frequent light cultivation.
4. Yields tend to be lower in older plantings because weeds and pests tend to build up over time.
5. There is variability in yield due to weather and variable pest pressure.

For a more comprehensive understanding of strawberry production we suggest the following resources: Strawberry Production Guide for the Northeast, Midwest, and Eastern Canada, NRAES-88 available for purchase from [Plant and Life Sciences Publishing](#) (PALS, formerly NRAES), and Strawberries: Organic Production, available for purchase from the [National Sustainable Agriculture Information Service, ATTRA](#). For those interested in strawberry production using day neutral strawberries we suggest: [Season-long Strawberry Production with Everbearers](#).

More research on growing perennial crops organically is needed, especially in the area of pest management. This guide attempts to compile the most current information available, but acknowledges that effective means of organic control are not available for some pests. Future revisions to this guide will incorporate new information providing organic growers with a complete set of useful practices to help them achieve success.

This guide uses the term Integrated Pest Management (IPM) which, like organic production, emphasizes the use of cultural practices to minimize pest outbreaks. With the limited pest control products available in many organic production systems, IPM techniques such as keeping accurate pest history records, selecting the proper site, and preventing pest outbreaks through use of sanitation, variety selection and biological controls are essential to producing a high quality crop.

All website addresses and links are listed in Section 11, References and Resources. A glossary of terms used in this guide is included at the end in section 12.

1. GENERAL ORGANIC MANAGEMENT PRACTICES

1.1 Organic Certification

Who needs to be certified?

Operations or portions of operations that produce or handle agricultural products that are intended to be sold, labeled, or represented as "100 percent organic," "organic," or "made with organic ingredients" or food group(s).

Farming operations that gross more than \$5,000 per year in organic products and want to use the organic label must be certified by a USDA National Organic Program (NOP) accredited certifying agency. The choice of certifier may be dictated by the processor or by the target market. A list of accredited certifiers operating in New York can be found on the New York State Department of Agriculture and Markets [Organic Farming Development/Assistance](#) web page. See more certification details in this guide under Section 3.1, Organic Certification Site Requirements.

Who does NOT need to be certified?

Producers and handling (processing) operations that sell less than \$5,000 a year in organic agricultural products do not need to be certified. Although exempt from certification, these producers and handlers must abide by the national standards for organic products and may label their products as organic. Handlers, including final retailers, that: do not process or repackage products; only handle products with less than 70 percent organic ingredients; process or prepare, on the premises of the establishment, raw and ready-to-eat food labeled organic; choose to use the word organic only on the information panel; and handle products that are packaged or otherwise enclosed in a container prior to being received by the operation and remain in the same package. More information can be found at the USDA Agricultural Marketing Service's [National Organic Program](#) (NOP) website.

1.2 Organic System Plan

An organic system plan (OSP) is central to the certification process. The OSP describes production, handling, and record-keeping systems, and demonstrates to certifiers an understanding of organic practices for a specific crop. The process of developing the plan can be very valuable in terms of anticipating potential issues and challenges, and fosters thinking of the farm as a whole system. Soil, nutrient, pest, and weed management are all interrelated on organic farms and must be managed in concert for success. Certifying organizations may be able to provide a template for the farm plan. The following description of the organic system plan is from the USDA [National Organic Program Handbook](#):

“A plan of management of an organic production or handling operation that has been agreed to by the producer or handler and the certifying agent and that includes written plans concerning all aspects of agricultural production or handling described in the Organic Food Production Act of 1990 and the regulations in [Subpart C](#), Organic Production and Handling Requirements.”

The [National Sustainable Agriculture Information Service](#), (formerly ATTRA), has produced a [Guide for Organic Crop Producers](#) that includes a chapter on writing the organic system plan. The [Rodale Institute](#) has also developed resources for transitioning to organic and developing an organic system plan.

It is important to note that the [USDA National Organic Program](#) requires that applicants for certification must keep accurate post-certification records for 5 years concerning the production, harvesting, and handling of agricultural products that are to be sold as organic. These records must document that the operation is in compliance with the regulations and verify the information provided to the certifying agent. Access to these records must be provided to authorized USDA representatives, including the certifying agent.

2. SOIL HEALTH

Healthy soil is the basis of organic farming. Regular additions of organic matter in the form of cover crops, compost, or manure create a soil that is biologically active, with good structure and capacity to hold nutrients and water. The minimum acceptable days to harvest interval for raw manure is 120 days (see National Organic Standards); buyers may require a period longer than 120 days between application and harvest however. Always maximize the time between the application of raw manure and harvest; in the case of perennial strawberry plantings, application during the planting year is recommended so that manure is not applied during a bearing year. It is important to never side dress with raw manure or use straw that has been used as animal bedding. Decomposing plant materials will support a diverse pool of microbes, including those that break down organic matter into plant-available nutrients as well as others that compete with plant pathogens in the soil and on the root surface. The practice of crop rotation to promote a healthy soil should be initiated in the one or two years prior to planting establishment. Organic growers must attend to the connection between soil, nutrients, pests, and weeds to succeed. An excellent resource for additional information on soils and soil health is *Building Soils for Better Crops*, 3rd edition, by Fred Magdoff and Harold Van Es, 2010, available from SARE, Sustainable Agriculture Research and Education, www.sare.org/publications/soils.htm. For more information, refer to Cornell's [Comprehensive Assessment of Soil Health website](#). In addition, a webinar series specifically about soil and nutrient management in berries is archived at: <http://www.fruit.cornell.edu/berry/production/soilnutrientmgmt/> along with *Berry Soil and Nutrient Management - A Guide for Educators and Growers*, <http://www.fruit.cornell.edu/berry/production/soilnutrientmgmt/pdfs/BerrySoilandNutrientManagementGuide.pdf>.

3. SITE SELECTION

For organic strawberry production, the importance of proper site selection and preparation cannot be over-emphasized. June-bearing strawberries are usually grown for two to three years in organic production systems, bearing fruit in the second and third years. Day neutral strawberries are usually grown for 2 years in organic production, bearing fruit in the first and second years. These approaches maximize yields while soil nitrogen content remains at acceptable levels. Consider that an ideal site should be close to your markets, be of sufficient acreage to allow for crop rotation, have available water of acceptable quality for irrigation and frost protection, have well-drained soil, and good air drainage (slopes of 3-4% preferably facing north and away from prevailing winter winds). Sites should not have recently been cropped to plants susceptible to Verticillium wilt.

Conduct needed site improvements prior to planting. Once strawberries are planted it is very difficult to make major changes to improve soil and air drainage, or to modify soil tilth, pH, or nutrient status. Improving soil structure or eliminating soil compaction layers in an established planting rarely prove feasible given the few years the crop is in the ground.

Weather plays a critical role in site selection, as well. The macroclimate, mesoclimate and microclimate of a strawberry site play important roles in variety selection and potential profitability. Of particular importance are the potential for spring frosts, winter minimum temperatures, length of the growing season, and growing season heat accumulation. More detailed information on the site selection information presented here also can be found in [Strawberry Production Guide for the Northeast, Midwest and Eastern Canada](#) (NRAES-88) available for purchase from [Plant and Life Sciences Publishing](#) (PALS, formerly NRAES).

A web-based, interactive site selection tool, the [New York Vineyard Site Evaluation System](#), uses specific climate information with a 3 kilometer resolution, based on 30 years of weather data, to determine the suitability of your site for different grape varieties. Although the tool was developed for vineyards, the map-based system integrates information on climate, topography, soils, and winter low temperatures much of which may be applicable to site selection for strawberry varieties across the state.

3.1 Organic Certification Site Requirements

The National Organic Program has requirements that affect site selection. Fields must not have been treated with prohibited products for three years prior to harvest of the certified organic crop. Mandated one-year crop rotation out of strawberries must be observed, though a 3-5 year rotation is typical. Adequate buffer zones must exist between certified organic and conventionally grown crops to prevent drift of prohibited materials onto certified organic crops. The buffer zones must be either a barrier (diversion ditch or dense hedgerow) or an area of sufficient size. The buffer zone needed will vary depending on equipment used on adjacent non-certified land. For example, use of high-pressure spray equipment or aerial pesticide applications in adjacent fields will increase the buffer zone size. Check with your certifier for specific buffer requirements. Buffer zone sizes commonly range from 20 to 250 ft, depending on adjacent field practices. Buffers can include windbreaks and living barriers such as a dense hedgerow. A dense hedgerow less than 50 ft wide may offer better protection from contamination than a 50-ft-wide open buffer zone. The [National Organic Farmers Association of New York \(NOFA NY\) organic certification guidance manual](#) states: "If the buffer is planted to the same crop as the field, documentation of what is done with the non-certified buffer crop is required. If harvested, non-certified harvest records and equipment decontamination logs should be maintained." Crops grown in the buffer zone may not be marketed as certified organic, or used for feed or bedding for certified organic livestock or dairy cattle.

3.2 Soil and Air Drainage and Soil Depth

Preparations for a strawberry planting should begin at least one year in advance. Selecting a site with good air and water drainage is essential for successful organic production. A nutritionally healthy planting in a well-drained soil with exposure to air movement is least susceptible to damage from pests and frosts.

Strawberries need good internal soil drainage to grow and do best on a well-drained sandy loam. Wet soils restrict root growth and respiration, resulting in weak growth and reduced yields. Coarse-textured soils have excellent soil drainage, but heavier soils, or soils with perched water tables often need drainage tiles to remove excess water and improve internal soil drainage. Drainage tile is best installed before planting. Local soil and water conservation districts and private tiling contractors can provide technical assistance in designing a drainage plan, but keep in mind that many base their designs on annual row crops. Perennial crops often require more intensive drainage than annual row crops. Planting on raised beds or on berms is useful to improve soil drainage in the rooting zone. Strawberries should not be grown on heavy clay soils. Because of the need for frequent light cultivation to manage weeds, stony and gravelly soils can also prove difficult.

Air drainage is an important consideration in choosing a strawberry field site. Cold air, like water, runs downhill, and collects in low areas or areas where trees or hedgerows obstruct airflow. These 'frost pockets' increase the risk of both mid-winter cold injury and spring frost damage. Selecting a site with a gentle slope (3-4%) and good air drainage will reduce the risk of cold or

frost injury. Good air drainage will also promote faster drying of foliage, flowers and fruit which will reduce the duration and frequency of disease infection periods. Good air drainage is essential to an organic disease management strategy.

Although strawberries can be grown on a wide variety of soils, shallow soils have less water holding capacity and will limit root development, resulting in smaller plants with smaller crops. Rooting depth of 12 inches or more is considered important for adequate plant growth and cropping levels. Digging test soil pits can help you evaluate potential rooting depth and drainage issues and evaluate what measures to take to address soil management issues before planting.

3.3 Soil Testing

Knowing all you can about the soil of a potential strawberry site will allow for better management decisions prior to planting. Soil testing is recommended to provide information on pH, availability of major and minor nutrients, organic matter and cation exchange capacity. A pH of 6.0 to 6.5 is suggested for most strawberry varieties. A [Cornell Soil Health Test](#) prior to planting will provide a comprehensive picture of soil condition, and includes nutrient analysis plus physical and biological analyses of the soil. See Table 6.1 for soil and tissue testing laboratories and refer to section 6, Nutrient Management, for more information.

A nematode analysis performed on representative soil samples is a wise step in the year or two prior to planting since it will allow time for using a cover crop to reduce plant parasitic nematode populations, see section 4, Cover Crops, for more information. Samples may be submitted for nematode testing to the Plant Disease Diagnostic Clinic, College of Agriculture and Life Sciences, Ithaca, NY. For more information and fee schedules visit their website at www.plantclinic.cornell.edu/. The best time for collecting samples for nematode testing is during summer, when soils are moist, not dry. A minimum of 6 soil subsamples, approx. 1" diameter and 4" deep should be collected randomly from an area approx. ½ acre in size. Gently mix samples together, transfer about 1 pint of mixed soil to a plastic bag, and ship as soon as possible to the diagnostic lab. Refrigerate sample if it cannot be shipped immediately.

3.4 Previous Cropping History

Another factor to consider when selecting a site is previous cropping history. The *Verticillium* wilt fungus may persist many years in soil and is devastating to strawberries under conditions favorable for disease development. If possible, avoid sites where potatoes, tomatoes, eggplants, or brambles have recently been grown and, to a lesser extent, squash, cucumber, pepper, or melons. These crops serve as hosts to *Verticillium* wilt. Many weeds are also hosts of the *Verticillium* fungus, particularly nightshade, groundcherry, redroot pigweed, lambsquarters, and horsetail. Weeds should be strictly controlled in current and future planting sites to keep *Verticillium* inoculum low. Rotating to non-susceptible grasses and cereals (5-8 year rotation) will reduce the amount of *Verticillium* inoculum in infested soil, but seldom eliminates it. Brassica crop rotations (mustards, broccoli, Brussels sprouts) are recommended where *Verticillium* wilt is present or has been observed in the past. Brassicas should be grown for a 2-yr period and crop residues incorporated into the soil. Practice long rotations out of strawberry and plant only resistant varieties where *Verticillium* wilt is a problem.

3.5 Irrigation Water Source

Another important criterion to consider when selecting a strawberry site is irrigation water quantity and quality. The irrigation water source should provide sufficient volume of water to irrigate as needed during the growing season. The irrigation system should be in place prior to planting to insure availability of water to the new transplants and to provide frost protection on cold nights during bloom. Trickle irrigation uses water more efficiently than overhead irrigation, but overhead irrigation can be used for frost protection. With trickle systems, row covers are required for frost protection in the absence of overhead irrigation. June-bearing strawberries, grown in a matted row system, typically require 1 to 2 inches of rainfall per week, or 25 to 30 inches per season. The critical periods when June-bearing strawberries require sufficient water to optimize growth and yield are during the fruiting development period through harvest and at renovation. Day neutral strawberries have similar water requirements but are likely to require more than twice the inches per season as June-bearing strawberries since they flower and fruit more or less continuously from mid-June to first frost in the fall.

Be sure to have a water test done on irrigation water sources prior to site selection to determine its physical, chemical, and biological constituents. Irrigation water pH should be 7.0 or below, and should also have a low salt content (<2.0 ds/m; preferably <1.0 ds/m) as strawberries are a salt-sensitive fruit crop. Always check with your certifier on the products used for lowering irrigation water pH. Water contaminated with sewage or manure should not be used to irrigate strawberries if it will come in contact with the berries. For more information on irrigation see the [Strawberry Production Guide for the Northeast, Midwest and Eastern Canada](#), available for purchase from [Plant and Life Sciences Publishing](#) (PALS, formerly NRAES).

4. COVER CROPS

Cover crops are grown for their valuable effect on soil properties, such as organic matter, and, in strawberries, on their preplant ability to eliminate or suppress weeds, provide nutrients to the plants, and reduce nematode populations. They can also improve water infiltration into the soil, maintain populations of beneficial fungi, and may help control insects and diseases. To be effective, cover crops should be treated as any other valuable crop on the farm, with their cultural requirements carefully considered and met, including nutrient requirements; susceptibility, tolerance, or antagonism to root pathogens and other pests; life cycle; and mowing/incorporation methods. See Table 4.1.1 for more information on specific cover crops.

4.1 Goals and Timing for Cover Crops

Cover crops play an important role in a strawberry planting, especially during the years prior to planting through improvement of soil organic matter, breaking up of compaction layers, erosion control, and suppression or elimination of weeds. Goals should be established for choosing a cover crop; for example, the crop can add nitrogen, smother weeds, or reduce nematode populations. The cover crop might best achieve some of these goals if it is in place for an entire growing season and incorporated into the soil prior to plant establishment.

Table 4.1.1. Cover Crops for Strawberries: Cultural Requirements and Crop Benefits

SPECIES	PLANTING DATES	LIFE CYCLE	SOIL TYPE PREFERENCE	SEEDING (LB/A)	COMMENTS
Alfalfa ¹	Early April-late May	Perennial	Well-drained, high pH (6.0-7.0)	14	+May be difficult to incorporate if allowed to overwinter
Brassicas e.g. mustards, rapeseed	April OR late Aug.-early Sept.	Annual /biennial	Loam to clay	5-12	+Good dual purpose cover & forage +Establishes quickly in cool weather +Mow or incorporate before seed formation +Biofumigant properties
Buckwheat	Late spring-early summer	Summer annual	Most	35-134	+Rapid grower (warm season) +Good catch or smother crop +Good short-term soil improver for poor soils +Mow or incorporate before seed formation +Will winter kill
Cereal Rye	August-early October	Winter annual	Sandy to clay loams	60-200	+Most cold-tolerant cover crop +Excellent allelopathic weed control +Good catch crop, rapid germination & growth +Mow or incorporate before seed formation +Temporary nitrogen (N) tie-up when turned under
Fescues fine (red, hard) tall	April-May OR late Aug.-Sept.	Long-lived perennial	Most	70-100	+Very good low-maintenance permanent cover, especially in infertile, acid, droughty &/or shady sites +Can be incorporated preplant +Tall fescue has high vigor, requires more frequent mowing, and has moderately high water use +Fine fescues have low vigor, require less frequent mowing, and have moderate water use
Marigold	Late May-June	Annual	Most	5-10	+Will winter kill +Biofumigant properties
Oats	Mid-April OR late Aug.-mid Sept.	Summer annual	Silt & clay loams	60-100	+Incorporate in late June when planted in the spring +Rapid growth +Ideal quick cover crop +When planted in late summer, will winter kill
Ryegrass	August-early Sept.	Winter annual OR short-lived perennial	Most	14-35	+Temporary N tie-up when turned under +Rapid growth +Good catch crop +Heavy N & moisture users
Sorghum-Sudangrass	Late Spring-Summer	Summer annual	NI	50-90	+Tremendous biomass producers in hot weather +Good catch or smother crop +Biofumigant properties
Sweet Clover ¹	Early April-mid May OR early August	Annual /biennial	Most	12-20	+Good dual purpose cover & forage +Does not need added nitrogen +May need to be mowed prior to incorporating +Mow or incorporate before seed formation
Vetch ¹	August	Annual /biennial	Most	30-40	+Does not need added nitrogen +Mow or incorporate before seed formation
Wheat	Early-mid Sept.	Winter annual	Most	80-100	+Mow or incorporate before seed formation

Adapted from M. Sarrantonio. 1994. Northeast Cover Crop Handbook; the Mid-Atlantic Berry Guide for Commercial Growers. 2008. Penn State Univ; the Pest Management Guidelines for Berry Crops. 2009. Cornell Univ.; and M. Pritts and D. Handley, eds. 1998. Strawberry Production Guide, NRAES-88.

¹ Legumes may benefit from inoculation of seed with nitrogen-fixing bacteria when planted in a field for the first time. Check with your certifier for allowable sources of inoculum.

Cover crops planted in late summer will suppress annual weed growth, improve soil texture, provide organic matter, and may increase soil nitrogen. The cover crop can be incorporated in late fall or in early spring before planting. Certain cover crops (marigold, sudangrass) will either suppress or resist nematode populations. In addition to producing large amounts of biomass that out-compete other plant species, some cover crops (annual rye, ryegrass) can inhibit weed growth through allelopathy, the chemical inhibition of one plant species by another. Rye provides allelopathic suppression of weeds when used as a cover crop, and when crop residues are retained as mulch. Rye residues retained on the soil surface release chemicals that inhibit germination and seedling growth of many grass and broadleaf weed species. Retention of residue on the soil surface can be accomplished by mowing before seed head formation.

See [Cornell's online cover crop decision tool](#) to match goals, season, and cover crop. Although written for vegetable growers it has comprehensive information on various cover crops. Another resource for determining the best cover crop for your situation is the Northeast Cover Crop Handbook, by Marianne Sarrantonio.

Allowing cover crop residue to remain on the soil surface might make it easier to fit into a crop rotation and will help to conserve soil water. Keep in mind that some of the nitrogen contained in the residue will be lost to the atmosphere, and total organic matter added to the soil will be reduced. Turning under the cover crop will speed up decomposition and nitrogen release from the crop residue. Cover crops such as grasses with low nitrogen content should be plowed under in the fall to allow time for decomposition prior to planting strawberries. Legumes which contain more nitrogen and decompose more quickly can be plowed under within a month of planting.

4.2 Legumes

Legumes are looked to as a potential nitrogen source. Legumes may benefit from inoculation of seed with nitrogen-fixing bacteria when planted in a field for the first time. Check with your certifier for allowable sources of inoculum. Legumes such as red clover and hairy vetch will often benefit from having a nurse crop planted simultaneously, usually a small cereal grain such as wheat or rye. These nurse crops establish faster than legumes and provide soil stability and reduce weed pressure during establishment, and provide support for the newly growing legumes before winter. To receive the full nitrogen benefit from planting legumes, they need to be incorporated into the soil just as they start to bloom, which is usually in late spring. (Source: Bjorkman, T. [Cover Crops for Vegetable Growers](#) website.)

5. VARIETY SELECTION

Key considerations in variety selection include the market destination and whether June-bearing or day neutrals will be grown. Consider whether the strawberries will be shipped and, if so, choose varieties with good shelf life and shipping quality. Flavor varies considerably among varieties and may be inversely related to shipping quality. Flavor may fluctuate depending on soil type, plant nutrition, and irrigation. Determine whether flavor or shipping quality are most important to your market and choose varieties accordingly. More information about strawberry varieties is available online, in the [Strawberry Production Guide for the Northeast, Midwest and Eastern Canada](#), available for purchase from [Plant and Life Sciences Publishing](#) (PALS, formerly NRAES), and in nursery catalogs.

In organic strawberry production, the variety's relative resistance or susceptibility to diseases is vital because of the limited number of organic fungicides that are available for disease management. June-bearing varieties considered to have the best potential for organic production in New York State include:

Earlyglow (early season), L'Amour (early/midseason), Jewel (midseason), Darselect (midseason), Winona (mid/late season), Allstar (mid/late season), Clancy (late season), Ovation (very late season).

Day neutral strawberry varieties considered to have the best potential for organic production in New York include Albion and Seascape.

Varieties vary widely in their susceptibility to fungal diseases and some may be less susceptible to insects. Jewel shows excellent fruit disease resistance but poor root disease resistance so raised beds are highly recommended. Darselect exhibits high root disease tolerance but may get powdery mildew in high fog or mist situations. If susceptible varieties are planted, the importance of site, sanitation and cultural practices will increase in accordance to the variety's susceptibility. Matching the variety to specific growing location can mean the difference between success and failure. Production practices such as raised beds, drip irrigation and plasticulture production may alleviate some pest pressure. Table 5.1 lists the relative disease susceptibility of many of the strawberry varieties grown in the Northeast. This is not an inclusive list and does not represent all varieties that are, or have been, grown organically in New York State.

Growers must also consider where they obtain their planting stock. According to language in the USDA-NOP regulation §205.202, “the producer must use organically grown seeds, annual seedlings, and planting stock. The producer may use untreated nonorganic seeds and planting stock when equivalent organic varieties are not commercially available. Seed and planting stock treated with substances that appear on the National List may be used when an organically produced or untreated variety is not commercially available. Planting stock used to produce a perennial crop may be sold as organically produced planting stock after it has been maintained under a system of organic management for at least 1 year. Seeds, annual seedlings, and planting stock treated with prohibited substances may be used to produce an organic crop when the application of the substance is a requirement of Federal or State phytosanitary regulations.” With the limited availability of organically certified strawberry stock, growers will likely be able to justify the use of non-organic stock to their certifying agency.

Table 5.1. Relative disease susceptibility among some strawberry varieties¹							
<i>Disease susceptibility^a</i>							
Variety	LSc	LSp	LB	RS^b	PM	VW	AT
Albion	U	I	U	R	I	R	S
Allstar	T-R	S-T-R	S	R-VR	T-R	I-T-R	VS
Annapolis	S	S	U	T-R	S	S	U
Cavendish	R	R	U	R	U	T-R	U
Chandler	U	S	S	S	R	U	VS
Clancy	T	T	T	R	R	R	R
Darselect	T	S	S	I	S	U	U
Earliglow	R	S-I-R	S	I-R	S-I-R	I-T-R	S
Evie II	U	U	U	T	S	T	U
Honeoye	T-R	S-T-R	U	S	S-I	S	U
Jewel	R	R	U	S	T	S	R
Kent	I-R	S-R	U	S	S	S	U
L'Amour	T	T	T	T	R	R	R
Lateglow	T-R	T-R	S	R	S	R-VR	U
Mesabi	T	T	U	R	U	R	U
Mira	U	U	U	R	U	U	U
Northeaster	T	T	U	R	U	R	U
Ozark Beauty	U	R	U	S	U	S	U
Redchief	R	S-R	VS	R	S-R	I-R	VS
Seascape	U	T	U	T	R	U	S
Sparkle	S-I	S-R	U	S-R	R	I-S	U
Tribute	T	T	U	R-VR	R	T-R	U
Tristar	T	T	U	R	R	R	U
AC Wendy	T	S	U	I	T	S	U
Winona	R	R	U	R	U	T	U

Key: VS = very susceptible, S = susceptible, I = intermediate, T = tolerant, R = resistant, VR = very resistant, U = unknown. Where multiple letter designations are given, ratings varied at different research sites.

¹The relative ratings in this chart apply to an average growing season. Under conditions favorable for disease development, any given variety may be more severely affected.

a. LSc=Leaf Scorch, LSp=Leaf Spot, LB=Leaf Blight, RS=Red Stele, BRR=Black Root Rot, PM=Powdery Mildew, VW=Verticillium Wilt, AT=Anthracnose.

b. Varieties are not resistant to all races of the red stele pathogen.

6. NUTRIENT MANAGEMENT

To produce a healthy crop, soluble nutrients must be available from the soil in amounts that meet the minimum requirements for the whole plant. The challenge in organic systems is balancing soil fertility to supply required plant nutrients at a time and at sufficient levels to support healthy plant growth. Restrictions in any one of the needed nutrients will slow growth and can reduce crop quality and yields. In strawberry plantings, the key considerations when managing nutrition organically include preplant soil

pH and nutrient adjustments; nutrition in established plantings; and understanding carbon to nitrogen ratios to deliver appropriate amounts of nitrogen to the crop.

Organic growers often speak of feeding the soil rather than feeding the plant. A more accurate statement is that organic growers focus their fertility program on feeding soil microorganisms rather than the plant. Soil microbes decompose organic matter to release nutrients and convert organic matter to more stable forms such as humus. This breakdown of soil organic matter occurs throughout the growing season, depending on soil temperatures, water availability and soil quality. The released nutrients are then held on soil particles or humus making them available to crops or cover crops for plant growth. Amending soils with compost, cover crops, or crop residues also provides a food source for soil microorganisms and when turned into the soil, starts the nutrient cycle again.

One goal of the grower is to heighten resource use efficiency (land, water, nutrients) to optimize plant growth and fruit yield. Plant size and yield can be influenced by water and nutrient supply (i.e. adequate water is needed for adequate nutrient uptake). Weak plants with few, small leaves will intercept insufficient sunlight to produce adequate yields in the current season or to develop flower buds for the next season. Conversely, over-stimulated plants with abundant large, dark green leaves have low water use efficiency, are more prone to winter injury, diseases and insect feeding, and produce fewer fruit. Organic strawberry plantings should strive to balance soil nutrient availability—via irrigation, organic matter content, soil pH, and microbial activity—with plant growth and production goals.

Nutrient demand for June-bearing strawberries is greatest during leaf and fruit development in spring when reserve nutrients carried over from the previous year have been used up and the plant is actively growing. Conversely, day neutral strawberries have a relatively consistent nutrient demand throughout the course of the season. Plant age, vegetative growth, and fruit yield are the deciding factors in determining the need for nutrients during the growing season.

A webinar series specifically about soil and nutrient management in berries is archived at:

<http://www.fruit.cornell.edu/berry/production/soilnutrientmgmt/> along with Berry Soil and Nutrient Management - A Guide for Educators and Growers, <http://www.fruit.cornell.edu/berry/production/soilnutrientmgmt/pdfs/BerrySoilandNutrientManagementGuide.pdf>.

6.1 Soil and Leaf Analysis

Regular soil and leaf analysis helps monitor nutrient levels. Choose a reputable nutrient testing lab (see Table 6.1.1) and use it consistently to avoid discrepancies caused by different extraction methods. It is recommended that annual leaf testing be incorporated into a fertility management program with biennial soil testing to assist in determining the plants' nutrient status and to make sure that what is in the soil is making it into the plants in the proper amounts. It is recommended that soil and leaf tests be completed in each block. Leaf testing is especially crucial in getting the information needed to make management decisions in problem areas of the planting and should be used on a more frequent basis, if needed.

Table 6.1.1. Nutrient Testing Laboratories

TESTING LABORATORY	WEB URL	SOIL	LEAF	COMPOST/MANURE	FORAGE
<i>Dairy One (Cornell Recommendations)</i>	http://dairystone.com/analytical-services/agronomy-services/about-agro-one/	x	x	x	x
<i>Agri Analysis, Inc.</i>	www.agrianalysis.com/		x	x	
<i>A&L Eastern Agricultural Laboratories, Inc.</i>	www.al-labs-eastern.com/	x	x	x	
<i>Cornell Soil Health (Cornell Recommendations)</i>	soilhealth.cals.cornell.edu/	x			
<i>Penn State Agricultural Analytical Services Lab.</i>	www.aasl.psu.edu/	x	x	x	
<i>University of Massachusetts</i>	http://www.umass.edu/soiltest/	x	x	x	
<i>University of Maine</i>	anlab.umesci.maine.edu/	x	x	x	x

Table 6.1.2. Deficient, sufficient, and excessive nutrient concentrations in strawberry leaves.

Nutrient	Symbol	<i>Target values (ppm, unless otherwise noted)</i>		
		Deficient Below	Sufficient	Excess Above
Nitrogen	N	1.90%	2.00-2.80%	4.00%
Phosphorus	P	0.20%	0.25-0.40%	0.50%
Potassium	K	1.30%	1.50-2.50%	3.50%
Calcium	Ca	0.50%	0.70-1.70%	2.00%
Magnesium	Mg	0.25%	0.30-0.50%	0.80%
Sulfur	S	0.35%	0.40-0.60%	0.80%
Boron	B	23	30-70	90
Iron	Fe	40	60-250	350
Manganese	Mn	35	50-200	350
Copper	Cu	3	6-20	30
Zinc	Zn	10	20-50	80

Adapted from: Pritts (1998) Soil and Nutrient Management. Chap. 7 In: Strawberry Production Guide. M. Pritts and D. Handley (eds.). NRAES-88. Ithaca, NY.

Note: ppm is parts per million.

% by dry weight of strawberry leaf

Table 6.1.2 gives the target values for strawberry leaf nutrients sampled in late July or early August in the Northeast. Regular soil testing helps monitor nutrient levels, in particular phosphorus (P) and potassium (K). The source of these nutrients depends on soil type and historic soil management. Some soils are naturally high in P and K, or have a history of manure applications that have resulted in elevated levels. Additional plant available nutrients are supplied by decomposed soil organic matter or through specific soluble nutrient amendments applied during the growing season in organically managed systems. Many types of organic fertilizers are available to supplement the nutrients supplied by the soil. ALWAYS check with your certifier before using any product to be sure it is approved.

6.2 Soil pH

Maintaining a soil pH range of 6.0 to 6.5 is recommended for strawberries. Use the soil test results to determine the appropriate amount of lime (raise pH) or sulfur (lower pH) to apply. The lime or sulfur requirement will depend on soil texture, current pH, and organic matter content. Follow the recommendations of the soil test and apply and incorporate sufficient lime or sulfur prior to planting. It typically takes one year for the applied lime or sulfur to raise or lower the soil pH, respectively. The slightly acid soil pH of 6.0 to 6.5 is required to help avoid micronutrient deficiencies.

Prilled sulfur formulations are preferred for soil application because they are easier to work with, provide better coverage, and are cheaper than powdered sulfur. Prilled sulfur takes about one year or more to oxidize and reduce soil pH; powdered sulfur takes 6 to 9 months. Likewise, finely ground lime is more difficult to work with, but it will raise the soil pH faster than coarse particles.

6.3 Managing Nutrients

Follow the recommendations of the soil test when adding nutrients to prepare a site for planting. Pay particular attention to the soil test results for potassium, phosphorus, magnesium, calcium, and boron. If interpreting your own soil tests, it is important to know the phosphorus extraction method used by your analytical lab in order to get a proper recommendation. When preplant recommendations are followed, additional potassium and phosphorus likely will not be required unless the soil is very sandy. However, potassium (K) demand by strawberry plants is relatively high, so make certain there is sufficient available potassium in the soil preplant. Boron is frequently low in fruit plantings throughout the Northeast. Note: Boron testing is not included in most standard soil test packages and should be selected as added test for strawberry soils.

Refer to **CALCULATING THE AMOUNT OF PESTICIDE TO USE** and Tables 9.1.1, 9.1.2, and 9.1.3 in Section 9.1 for converting amounts per acre to amounts needed for smaller areas and for measuring and mixing small amounts.

In established plantings, base fertilizer amounts on leaf analysis. In the event that potassium is required, a reasonable amount of potassium to apply, preferably in the fall, is up to 100 lb/acre. See table 6.3.1 for organic sources of potassium. Pay attention to the K/Mg ratio and if it is above 4, then additional magnesium should be applied with the potassium fertilizer to prevent inducing a magnesium deficiency: the K/Mg ratio should be less than 5.

Table 6.3.1. Available Potassium in Organic Fertilizers

Sources	Pounds of Fertilizer/Acre to Provide given Pounds of K ₂ O per acre:				
	20	40	60	80	100
Sul-Po-Mag 22% K ₂ O also contains 11% Mg	90	180	270	360	450
Wood ash (dry, fine, grey) 5% K ₂ O, also raises pH	400	800	1200	1600	2000
Alfalfa meal ^a 2% K ₂ O, also contains 2.5% N and 2% P	1000	2000	3000	4000	5000
Greensand or Granite dust 1% K ₂ O (x 4) ^b	8000	16000	24000	32000	40000
Potassium sulfate 50% K ₂ O	40	80	120	160	200

^a Only non-GMO sources of alfalfa may be used. Check with your certifier.

^b Application rates for some materials are multiplied to adjust for their slow to very slow release rates. Should be broadcast and incorporated prior to planting.

Table 6.3.2. Available Phosphorous in Organic Fertilizers

Sources	Pounds of Fertilizer/Acre to Provide given Pounds of P ₂ O ₅ Per Acre				
	20	40	60	80	100
Bone meal 15% P ₂ O ₅	130	270	400	530	670
Rock Phosphate 30% total P ₂ O ₅ (x4) ^a	270	530	800	1100	1300
Fish meal 6% P ₂ O ₅ (also 9% N)	330	670	1000	1330	1670

^a Application rates for some materials are multiplied to adjust for their slow to very slow release rates. Should be broadcast and incorporated prior to planting.

the excess nitrogen into the soil solution. When a material undergoing decomposition has an initially high C/N ratio (very little nitrogen), microbes will use whatever nitrogen is available for their own growth, leaving little for plants. This can result in temporary nitrogen deficiency. Once the decomposition process begins to slow and those microbes die off, they will release their nitrogen back into the soil where it will become available to plants. The rule of thumb is that if the C/N ratio is less than 20 or the material's nitrogen content is greater than 2.5%, then there will be enough nitrogen available for both decomposer microbes and plants. If the C/N ratio is above 20, then nitrogen will likely be immobilized until sufficient decomposition has taken place. One reason for applying nitrogen fertilizer at renovation is to help overcome the temporary nitrogen deficiency that will occur when the straw (with a high C/N ratio) is worked into the soil.

To create a robust organic fertility management plan, develop a plan for estimating the amount of nutrients that will be released from soil organic matter, cover crops, compost, and manure. A strategy for doing this is outlined in section 6.3. As these practices are integrated into field and farm management, the goal is to support diverse microbial communities that will help release nutrients from the organic matter additions. To assess overall impact of these practices on soil health, consider selecting a few target or problem fields for soil health monitoring over time via the [Cornell Standard Soil Health Analysis Package](#). This suite of eight tests complements a standard soil chemical nutrient analysis by focusing on biological and physical soil health indicators. While the test results will provide feedback on how the soil sample compares to other New York soils, the real power is in the baseline readings for comparison in the future after implementing new soil health management strategies.

Included in the Soil Health Test is an analysis of soil protein content. As with the other soil health tests, this serves as an indicator of soil management and amendment history. The test measures organic soil N that is in the form of proteins- an important food source for soil microbes. Use this test to help monitor impact and target future investments of legume cover crops and compost / manure applications.

Magnesium (Mg) deficiency in strawberry is quite common. Factors that influence magnesium availability include soil pH and excess potassium. In established plantings that are low to deficient in magnesium, typical recommendations would be for 10-40 lb/acre actual magnesium, but follow recommendations of the leaf analysis.

Boron is frequently low in fruit plantings throughout the Northeast. If boron is required, then apply no more than 2 lb/acre actual boron in any one year. The best time to apply boron is after leaves are mowed at renovation. Check with your certifier for information on allowable sources of magnesium and boron.

Phosphorus demand by strawberry is relatively low, and phosphorus is usually not required in established plantings. Table 6.3.2 lists some organic fertilizer sources of P.

6.4 Preparing a Nitrogen Budget

The carbon to nitrogen (C/N) ratio in compost can provide a guide for nitrogen release into the soil solution. When a decomposing material has a low C/N ratio (a lot of nitrogen) microbes release

Management of N, and insuring adequate supply at the times of crop need, requires some planning. Prepare a nitrogen budget for organic production to estimate the amount of N released by various organic amendments as well as native soil organic matter. Examples of manures and their nutrient content are shown in Table 6.4.1. Compost and manure should be tested for nutrient content at an analytical lab, and cover crops can be tested at a forage testing lab (Table 6.1.1). Knowing these values will help evaluate if the budget plan is providing appropriate amounts of N during the season by comparing them to the nitrogen guidelines for strawberries (Table 6.4.2)

Table 6.4.1. Estimated Nutrient Content of Common Animal Manures

	N	P ₂ O ₅	K ₂ O	N1 ¹	N2 ²	P ₂ O ₅	K ₂ O
	NUTRIENT CONTENT LB/TON			AVAILABLE NUTRIENTS LB/TON IN FIRST SEASON			
Dairy (with bedding)	9	4	10	6	2	3	9
Horse (with bedding)	14	4	14	6	3	3	13
Poultry (with litter)	56	45	34	45	16	36	31
Compost (from dairy manure)	12	12	26	3	2	10	23
Composted poultry manure	17	39	23	6	5	31	21
Pelletized poultry manure ³	80	104	48	40	40	83	43
Swine (no bedding)	10	9	8	8	3	7	7
	NUTRIENT CONTENT LB/1000 GAL.			AVAILABLE NUTRIENTS LB/1000 GAL FIRST SEASON			
Swine finishing (liquid)	50	55	25	25*	20+	44	23
Dairy (liquid)	28	13	25	14*	11+	10	23

1-N1 is the total N available for plant uptake when manure is incorporated within 12 hours of application.

2-N2 is the total N available for plant uptake when manure is incorporated after 7 days.

3 –Pelletized poultry manure compost. Available in New York from Kreher's.

* injected, + incorporated.

Adapted from "Using Manure and Compost as Nutrient Sources for Fruit and Vegetable Crops" by Carl Rosen and Peter Bierman and Penn State Agronomy Guide 2007-8.

Using the values from your soil test, estimate that 20 lbs. of nitrogen will be released from each percent organic matter in the soil. From the test of total N in any manure applied, estimate that 50% is available in the first year, and then 50% of the remaining is released in each of the next two years. So, for an application rate of 100 lbs. of N as manure, 50 lbs. would be available the first year, 25 lbs. the second, and 12.5 lbs. the third. Remember to check with your certifier on the days-to-harvest interval when using raw manure and allow a minimum of 120 days between application and harvesting. To prevent run-off, do not apply raw manure to bare ground in established strawberry plantings.

Estimate that between 10% and 25% of the N contained in compost will be available the first year. It is important to test each new mix of compost for actual amounts of the different nutrients available. Compost maturity will influence how much N is available. If the material is immature, more of the N may be available to the crop in the first year. A word of caution: Using compost to provide for a crop's nutrient needs is not generally a financially viable strategy. The total volume, trucking, and application can be very expensive for the units of N available to the crop. Most stable composts should be considered as soil conditioners, improving soil health, microbial diversity, tilth, and nutrient retaining capacity.

Add together the various N values from these different organic sources to estimate the N supplying potential of the soil. There is no guarantee that these amounts will actually be available in the season, since soil temperatures, water, and crop physiology all impact the release and uptake of these soil nutrients. If early in the organic transition, a grower may consider increasing the N budget supply by 25%, to help reduce some of the risk of N being limiting to the crop. Remember that with a long-term approach to organic soil fertility, the N mineralization rates of the soil will increase. This means that more N will be available from organic amendments because of increased soil microbial activity and diversity. Feeding these organisms different types of organic matter is essential to help build this type of diverse biological community and ensure long-term organic soil and crop productivity.

Table 6.4.2. Annual Nitrogen Guidelines for June-bearing strawberries.

Planting Age (years)	Amount Actual N (lbs/Acre)	Time of Year to Apply
0	30	early June ^a
	30	early Sept ^a
1+	70	at renovation
	30	early Sept ^b

^a Be sure plants are growing well prior to application.

^b Adjust amount based on leaf analysis.

May and early June. Be aware that spring applications of nitrogen can greatly increase the risk of gray mold fruit rot infections.

Day neutral strawberries generally require 3 lb actual nitrogen per week during active spring growth. This rate should be increased to 5 lb actual N per week when fruiting begins. During the fruiting period, plants may require additional potassium; alternate a nitrogen fertilizer every other week with a nitrogen fertilizer product that supplies both N and K.

Table 6.4.3 lists some commonly available fertilizers, their nutrient content, and the amount needed to provide different amounts of available nitrogen, adapted by Vern Grubinger from the [University of Maine soil testing lab](#).

Table 6.4.3. Available Nitrogen in Organic Fertilizers

Sources	Pounds of Fertilizer/Acre to Provide given Pounds of N per Acre				
	20	40	60	80	100
Blood meal 13% N	150	310	460	620	770
Soy meal 6% N (x 1.5) ^a , also contains 2% P and 3% K ₂ O	500	1000	1500	2000	2500
Fish meal 9% N, also contains 6% P ₂ O ₅	220	440	670	890	1100
Alfalfa meal 2.5% N also contains 2% P and 2% K ₂ O	800	1600	2400	3200	4000
Feather meal 15% N (x 1.5) ^a	200	400	600	800	1000

^a Application rates for some materials are multiplied to adjust for their slow to very slow release rates.

7. ORGANIC STRAWBERRY IPM

Organic production of strawberries is challenging in New York State given the abundant rainfall during the growing season leading to increased pressure from diseases, insects and weeds. However, growers in New York and the eastern United States, through proper variety and site selection, strict attention to cultural practices and sanitation, and increased attention paid to scouting plantings on a weekly basis to catch pest outbreaks early, have succeeded in producing quality organic strawberries. In contrast, a failure to appreciate the risk of disease, insect and weed development, and failure to devise and implement a season-long (and multiyear) management strategy, can lead to serious crop and even plant losses in particular years. Successful IPM is essential to the sustainable production of organic strawberries.

7.1 Developing a Strawberry IPM Strategy

1. Examine your strawberry operation closely. Break it down into specific plantings, or “strawberry blocks.”
2. Produce a map of each planting (or block) to record weeds, pest outbreaks, nutrient deficiencies, drainage problems, missing plants, and any other abnormalities you find.
3. Develop a record-keeping system for each planting or block.
4. Develop a scouting plan for each block and record results.

5. Monitor and record weather factors and understand basic weather patterns of the area.
6. Keep accurate records of spray applications, tools, or tactics used to manage pests.
7. Properly maintain your spray equipment, calibrate the sprayer, select appropriate nozzles, and reduce spray drift. Consult the [Pesticide Application Technology website](#) at Cornell University or the [Strawberry Production Guide for the Northeast, Midwest, and Eastern Canada](#) (NRAES-88) available for purchase from [PALS Publishing](#).
8. Develop a thorough knowledge of the strawberry pests you are likely to encounter during the year. This includes basic pest biology, symptoms or damage, whether they are a primary or secondary pest, scouting thresholds, and the best time to implement management practices.
9. Choose a pest management strategy for the planting (or block) that is based on all of the information you've gathered. Use the options that make the most sense for your operation.
10. Continue your pest management education.

Other resources available online, include:

New York State IPM website: nysipm.cornell.edu/fruits/

Cornell Fruit Resources: www.fruit.cornell.edu/berry/

New York State berry IPM insect and disease fact sheet index: nysipm.cornell.edu/factsheets/berries/

Cornell University Pesticide Management Education Program: pmep.cce.cornell.edu/

Pesticide Application Technology at Cornell University: web.entomology.cornell.edu/landers/pestapp/

Elements of IPM for Strawberries in New York State www.nysipm.cornell.edu/elements/strawb.asp

Network for Environment and Weather Applications (NEWA) newa.cornell.edu

Berry Diagnostic Tool www.fruit.cornell.edu/berrytool/

7.2 Weed Management

Weed management is a major challenge for strawberry growers. Weeds are part of the strawberry planting ecosystem and can compete for water and nutrients; provide alternate hosts for pests; and interfere with planting operations. Weed growth can also alter the microclimate around plants, leading to higher disease pressure. In organic production, site preparation prior to planting spanning 2- to 3-years to eliminate weeds through cover cropping and cultivation will provide lasting benefits in weed control for the short-term perennial production cycle of strawberries. Table 7.2.1 outlines weed management practices in strawberry plantings.

Table 7.2.1. Weed management without herbicides in a strawberry planting.

Year	Month	Non-herbicidal options
Planting year ¹ ¹CRITICAL TIME FOR REDUCING WEEDS.	April - May	Till to prepare for planting.
	May	Cultivate.
	Mid-June after planting	Cultivate.
	Mid-July	Cultivate.
	Mid-August	Cultivate.
	October	Cultivate.
	Late November	Mulch for winter protection.
Fruiting years	March - April	Remove mulch.
	Early May	Hand weed only.
	Late July after harvest	Mow leaves, narrow rows with a tiller.
	September	Cultivate.
	November	Mulch for winter protection.

Excellent preplant preparation with the goal of eliminating perennial weeds from the site before planting is essential. Good preplant preparation, use of cover crops, and crop rotation help reduce weed pressure considerably. Eliminating perennial weeds can be achieved with repeated cultivation and using “green manure” cover crops that are plowed under prior to planting. For more information on cover crops see section 4. Keep in mind that excessive cultivation can lead to undesirable consequences such as soil erosion, reduced soil organic matter, and breakdown in soil structure resulting in compaction and reduced permeability.

Minimizing weed competition during plant establishment is critical to achieve optimal plant growth and yields. Once plants are set, regular hand weeding, hoeing, and cultivation are required throughout the first year. Do not let weeds go to seed, and keep the surrounding area mowed to prevent weed seeds from migrating into the planting site. If a first year planting is healthy, dense, and weed free prior to winter weed problems will be much less in subsequent years. Some growers are planting in late May or early June at a higher density to reduce weed pressure.

Managing weeds within the row may be one of the most difficult tasks in the production of organic strawberries. Inorganic mulches like plastic can only be used in organic production if they are removed from the soil annually. There has been some recent research in Italy with the use of biodegradable mulch films (starch-based) that do not need to be removed from the soil. These materials have shown promise in New York strawberry plantings.

Organic mulches can also be used as tools for weed management. They are most effective where soil moisture and fertility are low and where low plant size restricts crop productivity. To provide adequate weed control, organic mulches must be at least 4 inches thick. Potential organic mulches include straw, hay, sawdust, and wood chips. Mulch matted row plantings with straw (wheat or rye works best) for winter protection, then rake the straw into the alleyways for additional weed suppression. Straw mulch may serve as a major source of weed seed; be sure to inspect straw before purchase. Use of straw or hay mulch between the rows for suppression of weed growth is also an excellent method of water conservation and increasing the soil organic matter. Financial assistance to help pay for mulch may be available from your county's Soil and Water Conservation District office.

There are a number of mechanical, thermal and animal measures that can be used to limit the effects of weeds in a strawberry planting. Mechanical and thermal options include fixed hoes, rotary cultivators, flammers, steamers, and hot water applicators. Animal weeders have also been used with some success in organic plantings across the United States. The use of weeder geese, guinea fowl, and sheep have some effectiveness, but due to food safety concerns regarding microbial contamination of food crops from manure they should be used during the planting (non-bearing) year only. The mechanical brush hoe, in particular, showed promise for use in matted row strawberry production. Just two well-timed passes provided excellent seasonal weed control. The brushes moved runners back into the row, allowing cultivation to occur later in the season compared with other implements. The resulting layer of dust created by the implement "mulched" the field and suppressed weed seed germination.

Herbicides are applied in a dilute spray in a 4 foot strip under the plant. Consult the pesticide label for specifics on how to apply the pesticide. Note that you may need to use up to 100 gallons of solution per sprayed acre.

Note: An organic herbicide strategy alone cannot provide satisfactory weed control for organic strawberry growers.

At the time this guide was produced, the following materials were available in New York State for managing weeds and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Organic Herbicides Labeled for Management of Weeds in Strawberry					
Trade Name (<i>active ingredient</i>)	Product Rate	PHI (days)	REI (hours)	Efficacy ¹	Comments
AXXE Broad Spectrum Herbicide (<i>ammonium nonanoate</i>)	6-15% solution	-	24	?	Apply spray until the undesirable plants are fully wetted with the herbicide solution.

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?- not reviewed or no research available.

PHI - pre-harvest interval, REI - re-entry interval, - = pre-harvest interval isn't specified on label.

7.3 Principles of Insect and Disease Management

While strawberry production may be severely limited by insect pests and plant diseases, an understanding of the factors involved in their development can ensure effective management. The development of disease and insect damage is highly dependent on characteristics and conditions of the crop (host), the pathogen/pest population, and the environment. These factors all must be conducive before disease development and/or considerable insect damage will occur.

Characteristics of the host that influence disease and pest susceptibility include the host's vigor, physiology, and variety (genetics). Aggressiveness or virulence, abundance, and physiology are characteristics of the pest or pathogen populations that influence their ability to cause disease or damage. At the same time, abiotic environmental conditions such as temperature, moisture, light, and soil chemistry can affect both the host and pest and may promote or prevent disease. Moreover, the presence, abundance and activity of natural enemies can play an important role in determining pest status. The most successful disease pathogens and insect pests have coevolved with their hosts over many years to infect disease and damage at the most opportune times. To successfully minimize disease and pest damage, the relevant aspects of the host, pathogen/pest, and environment must all be managed within specific timeframes.

Although insect pests and plant disease pathogens are vastly different in their biology, they often have enough similarity in life history strategies to allow successful management under a single set of underlying principles. These principles include avoidance/exclusion, eradication, and protection. They are defined below.

Avoidance/exclusion

This principle focuses on preventing pathogen introduction and minimizing factors that favor the establishment of pests and pathogens. Several practices that exclude or limit pathogen and pest presence include the following:

- Select sites with good soil drainage. Install tile in plantings with less than optimal drainage and/or incorporate raised beds or berms to further promote soil drainage.
- Choose sites with good air drainage. Promote air circulation by selecting an open site, removing dead or senescent plant material and reducing weeds; these practices allow fruit and leaves in berry plantings to dry more quickly.
- Plant only disease free and insect free planting stock.
- Prevent rain-splash dispersal of soil particles by applying a thick layer of mulch under and around plants.
- Practice weed management as weeds can be hosts for strawberry pathogens and arthropod (insect and mite) pests.
- Avoid planting strawberries in proximity to other crops or habitats that harbor large pathogen and/or pest populations.
- Plant strawberries under covered production to avoid outside sources of disease inoculum.

Eradication

This principle is concerned with the destruction of pathogen/pest populations. These practices include:

- Sanitation of plantings by removal of infected/infested plant material including overripe fruit, leaf litter, and plants to eradicate pathogen and pest populations. Destruction of this material is accomplished through burning, chipping, burying, and composting.
- Several biological control alternatives are available for insect suppression for strawberry crops including products based on formulated *Bacillus thuringiensis* and insectary-reared predatory mites. Currently, there are no consistently reliable biological control products that have been developed for managing strawberry diseases, although there are numerous biopesticides that are available and effective in low disease pressure situations.
- Chemical application of fungicides, insecticides, and miticides may reduce pathogen and pest populations below damage thresholds, but will rarely eradicate them.

Protection

This principle is founded on protection of plants from pathogen infection and pest damage. Practices that protect plants by minimizing factors favoring infection and damage include the following:

- Plant strawberry varieties that are disease resistant or less susceptible to diseases of concern.
- Consider the use of protected production structures such as low tunnels, to reduce occurrence of fruit rots.
- Avoid excessive nitrogen fertilization as many pathogens, insects and mites thrive on succulent tissues.
- Keep fruit from contacting soil by use of mulch under and around the plants.
- Harvest fruit promptly and cool it to protect from fruit rots and insect infestations on overripe fruit.
- Applications of fungicides, insecticides, or miticides may protect susceptible tissues from disease and insect damage.
- Plant strawberries under covered production to protect against rainfall, which is the primary factor driving infection and spread of disease.

7.4 DISEASES OF PRIMARY CONCERN

Several important diseases that occur in the temperate climate of the northeastern U.S. are described below to help growers manage them with appropriate organic practices.

7.4.1 Leaf Blight (*Phomopsis obscurans*)

Leaf lesions begin as small, circular to irregular, reddish, or purplish spots. As they expand, lesion centers become necrotic and turn light brown with a dark purple halo. Older lesions along major leaf veins develop into large V-shaped lesions that eventually kill the leaf. Heavy leaf infections can inhibit the production of flower buds for the following year, predispose a plant to winter injury, and provide inoculum for infection of the fruit caps. Fruit may also be infected in some instances.

Leaf Blight Management Options	
Scouting/thresholds	None established.
Variety susceptibility	There are no reports of cultivar resistance to leaf blight but Jewel shows low infection rates
Cultural management	Destroying infected leaves at renovation (e.g., mowing and burying) will reduce the amount of carry-over inoculum. Promoting air circulation (plant spacing and weed control) will reduce foliage drying time and limit infection periods.
Chemical treatment	An early season fungicide application is recommended when carry-over inoculum from the previous year is high or conditions are favorable for disease development.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.4.1 Pesticides Labeled for Management of Leaf Blight					
Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Agricure (potassium bicarbonate)	2-5 lb/acre	0	1	?	
Badge X2 (copper hydroxide, copper oxychloride)	1-2.5 lb/acre	0	48	1	
Champ WG (copper hydroxide)	2-3 lb/acre	-	48	1	May cause crop injury under some conditions.
CS 2005 (copper sulfate pentahydrate)	19.2-25.6 oz/acre	0	48	1	
Cueva Fungicide Concentrate (copper octanoate)	0.5-2 gal/acre	up to day	4	1	
Milstop (potassium bicarbonate)	2-5 lb/acre	0	1	?	Do not mix with other pesticides or fertilizers. Not compatible with alkaline solutions.
Nu-Cop 50 WP (copper hydroxide)	2-3 lb/acre	1	24	1	Use higher rate when conditions favor disease. Discontinue use if signs of phytotoxicity appear. Copper may cause blue spotting on fruit.

Table 7.4.1 Pesticides Labeled for Management of Leaf Blight

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Nu-Cop 50DF (copper hydroxide)	2-3 lb/acre	1	48	1	See comment for Nu-Cop 50 WP.
Nu-Cop HB (copper hydroxide)	1-1.5 lb/acre	1	48	1	Discontinue use if signs of phytotoxicity appear.
Oxdate 2.0 (hydrogen dioxide, peroxyacetic acid)	32 fl oz – 1 gal/100 gal water	0	until dry	?	At-planting and existing planting foliar application.
PERpose Plus (hydrogen peroxide)	1 fl oz/gal Initial/curative	-	until dry	?	For initial or curative use, apply higher rate for 1 to 3 consecutive days. Then follow with weekly/preventative treatment.
PERpose Plus (hydrogen peroxide)	0.25-33 fl oz/gal Weekly/preventative foliar spray	-	until dry	?	For weekly or preventative treatments, apply lower rate every five to seven days. At first signs of disease, use curative rate then resume weekly preventative treatment.
Regalia (<i>Reynoutria sachalinensis</i>)	1-3 qt/acre	0	4	?	Initiate at first sign of disease then every 7-14 days.
Trilogy (neem oil)	1% solution	up to day	4	?	Maximum labeled use of 2 gal/acre/application

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.

PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.4.2 Leaf Scorch (*Diplocarpon earliana*)

Dark purple leaf spots about one eighth to one quarter inch in diameter appear scattered over the upper leaf surfaces or petioles. These spots differ from those of leaf spot in that they are purple throughout (no light centers). Numerous infections can cause a leaf to appear red or light purple and eventually to dry up and appear scorched. Heavy leaf infections can inhibit the production of flower buds for the following year, predispose a plant to winter injury, and provide inoculum for infection of the fruit caps.

Leaf Scorch Management Options

Scouting/thresholds	None established.
Variety susceptibility	Resistance and tolerance has been reported for several varieties. However, reports from different states often conflict; hence resistance/tolerance may be variable and/or region dependent. A consensus of reports suggests that 'Allstar', 'Jewel', 'Cavendish', 'Earliglow', 'Lester', and 'Redchief' have some resistance. 'Tristar' and 'Tribute' are susceptible but tolerant of infection.
Cultural management	Destroying infected leaves at renovation (e.g., mowing and burying) will reduce the amount of carry-over inoculum. Promoting air circulation (plant spacing and weed control) will reduce foliage drying time and limit infection periods.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.4.2 Pesticides Labeled for Management of Leaf Scorch

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Badge X2 (copper hydroxide, copper oxychloride)	1-2.5 lb/acre	0	48	1	
CS 2005 (copper sulfate pentahydrate)	19.2-25.6 oz/acre	0	48	1	
Cueva Fungicide Concentrate (copper octanoate)	0.5-2 gal/acre	up to day	4	1	
Serifel (Bacillus amyloliquefaciens)	4-16 oz/acre	0	4	?	

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.

PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.4.3 Leaf Spot (*Mycosphaerella fragariae*)

Initial lesions on leaves begin as small, irregularly shaped purple spots. Mature lesions become approximately one eighth to one quarter inch in diameter; remain relatively round, and the centers of lesions turn from a purplish brown to grayish white. The pathogen primarily infects young, expanding leaves and petioles, and occasionally fruit (black seed). Heavy leaf infections can inhibit the production of flower buds for the following year, predispose a plant to winter injury, and provide inoculum for infection of the fruit caps.

Leaf Spot Management Options

Scouting/thresholds	None established.
Variety susceptibility	Resistance and tolerance has been reported for several varieties. However, reports from different states are often in conflict with one another; hence resistance/tolerance may be variable and/or region dependent. A consensus of reports suggests that 'Jewel', 'Cavendish', 'Winona', and 'Lester', have some resistance. 'Tristar' and 'Tribute' are susceptible but tolerant of infection.
Cultural management	Destroying infected leaves at renovation (e.g., mowing and burying) will reduce the amount of carry-over inoculum. Promoting air circulation (plant spacing and weed control) will decrease foliage drying time and limit infection periods.
Chemical treatment	An early season fungicide application is recommended when carry-over inoculum from the previous year is high or conditions are favorable for disease development.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.4.3 Pesticides Labeled for Management of Leaf Spot					
Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy¹	Comments
Badge X2 (copper hydroxide, copper oxychloride)	1-2.5 lb/acre	0	48	1	
Basic Copper 53 (basic copper sulfate)	2-2.8 lb/acre	up to day	48	1	Copper may cause blue spotting on fruit.
Champ WG (copper hydroxide)	2-3 lb/acre	-	48	1	May cause crop injury under some conditions.
CS 2005 (copper sulfate pentahydrate)	19.2-25.6 oz/acre	0	48	1	
Cueva Fungicide Concentrate (copper octanoate)	0.5-2 gal/acre	up to day	4	1	
Nordox 75 WG (cuprous oxide)	3-5 lb/acre			1	Begin application when plants are established and then on a weekly basis.
Nu-Cop 50 WP (copper hydroxide)	2-3 lb/acre	1	24	1	Use higher rate when conditions favor disease. Discontinue use if signs of phytotoxicity appear. Copper may cause blue spotting on fruit.
Nu-Cop HB (copper hydroxide)	1-1.5 lb/acre	1	48	1	Discontinue use if signs of phytotoxicity appear.
PERpose Plus (hydrogen peroxide)	1 fl oz/gal Initial/curative 0.25-33 fl oz/gal Weekly/preventative	-	until dry	?	For initial or curative use, apply higher rate for 1 to 3 consecutive days. Then follow with weekly/preventative treatment. For weekly or preventative treatments, apply lower rate every five to seven days. At first signs of disease, use curative rate then resume weekly preventative treatment.
Regalia (<i>Reynoutria sachalinensis</i>)	1-3 qt/acre	0	4	?	Initiate at first sign of disease then every 7-14 days.
Serifel (<i>Bacillus amyloliquefaciens</i>)	4-16 oz/acre	0	4	?	
Trilogy (neem oil)	1% solution	up to day	4	?	Maximum labeled use of 2 gal/acre/application

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.

PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.4.4 Powdery Mildew (*Podosphaera aphanis*)

The edges of infected leaves roll up, sometimes revealing a white, powdery layer of mycelium and spores on the lower leaf surfaces. Purple to reddish blotches also occur frequently on the lower leaf surfaces. Symptoms are usually not evident until middle or late summer. Numerous pepper-like black flecks (overwintering spore-producing structures – cleistothecia) may appear on infected leaf surfaces in fall.

Powdery Mildew Management Options	
Scouting/thresholds	None established.
Variety susceptibility	Varieties such as 'Chandler', 'Clancy', 'L'Amour', 'Jewel', 'Tribute' and 'Tristar' rarely if ever show infection. Few varieties show infection in open field conditions until after harvest with the exception of 'Darselect' in high fog or mist locations. Leaf infections after renovation of 'Earliglow', 'Darselect', 'Evangeline', 'Annapolis', 'Raritan', 'Flavorfest', and 'Ovation' are common but usually do not cause economic damage.
Cultural management	Manage weeds and regulate planting density to promote good air circulation. Avoid excessive nitrogen and sites with poor air drainage. This disease may be more problematic in covered production where air circulation is reduced. High and low tunnel production can exacerbate powdery mildew problems.
Chemical treatment	See table below.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.4.4 Pesticides Labeled for Management of Powdery Mildew					
Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Acoidal (sulfur)	5-10 lb/acre	-	24	1	Begin applications when disease first appears. Repeat as necessary. Do not use on sulfur sensitive varieties.
Actinovate AG (<i>Streptomyces lydicus</i> WYEC 108)	3-12 oz/acre	0	until dry	3	For best results apply with a spreader/sticker prior to onset of disease. Re-apply at 7-14 day intervals depending on disease pressure and environmental conditions.
Agricure (potassium bicarbonate)	2-5 lb/acre	0	1	1	
Cinnerate (cinnamon oil)	13-30 fl.oz./100 gal water	-	-	?	25(b) pesticide.
Cueva Fungicide Concentrate (copper octanoate)	0.5-2 gal/acre	up to day	4	1	
Defend DF (sulfur)	5-10 lb/acre	-	24	1	Begin applications when disease first appears. Repeat as necessary. Do not use on sulfur sensitive varieties.
Double Nickel 55 (<i>Bacillus amyloliquefaciens</i> str D747)	0.25-3 lb/acre	0	4	2	
Double Nickel LC (<i>Bacillus amyloliquefaciens</i> str D747)	0.5-6 qt/acre	0	4	2	
Glacial Spray Fluid (mineral oil)	0.75 gal/100 gal	up to day	4	1	See label for specific application volumes and equipment.
GrasRoots (cinnamon oil)	1 part GrasRoots:9	-	-	?	25(b) pesticide.

Table 7.4.4 Pesticides Labeled for Management of Powdery Mildew

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
	parts water				
JMS Stylet-Oil (paraffinic oil)	3 qt/100 gal water	0	4	2	A high volume of water is needed for thorough coverage. Many common pesticides are phytotoxic when applied with or close to oil sprays (e.g., sulfur). Check label for restrictions.
Kaligreen (potassium bicarbonate)	2.5-3 lb/acre	1	4	3	Do not mix with highly acidic products or nutrients.
Micro Sulf (sulfur)	5-10 lb/acre	-	24	1	Some varieties may be sensitive to sulfur.
Microthiol Disperss (sulfur)	5-10 lb/acre	-	24	1	Not recommended within 2 weeks of an oil application nor if temperatures are expected to exceed 90 degrees within 3 days following the application.
Mildew Cure (garlic oil, cottonseed oil, corn oil)	1 gal/100 gal water acre	-	-	?	25(b) pesticide. Conduct phytotoxicity test prior application.
Milstop (potassium bicarbonate)	2-5 lb/acre	0	1	3	Do not mix with other pesticides or fertilizers. Not compatible with alkaline solutions.
M-Pede (insecticidal soap)	1-2% vol/vol	0	12	?	Curative control.
Organocide (sesame oil)	1-2 gal/100 gal water /acre	-	-	?	25(b) pesticide.
Oxidate 2.0 (hydrogen dioxide, peroxyacetic acid)	64 fl oz/100 gal water dip	0	until dry	1	Pre-plant dip.
Oxidate 2.0 (hydrogen dioxide, peroxyacetic acid)	32 fl oz – 1 gal/100 gal water	0	until dry	1	At-planting foliar application and foliar and crown disease control for existing plantings.
PERpose Plus (hydrogen peroxide)	1 fl oz/gal Initial/curative	-	until dry	3	For initial or curative use, apply higher rate for 1 to 3 consecutive days. Then follow with weekly/preventative treatment.
PERpose Plus (hydrogen peroxide)	0.25-33 fl oz/gal Weekly/pre-ventative	-	until dry	3	For weekly or preventative treatments, apply lower rate every five to seven days. At first signs of disease, use curative rate then resume weekly preventative treatment.
PureSpray Green (white mineral oil)	0.75-1.5 gal/100 gal water	up to day	4	?	Spray at no less than 400 PSI using ceramic nozzles.
Serifel (<i>Bacillus amyloliquefaciens</i>)	4-16 oz/acre	0	4	?	
Sil-Matrix (potassium silicate)	0.5-1% solution.	0	4	?	
SuffOil-X (aliphatic petroleum solvent)	1-2 gal/100 gal water /acre	up to day	4	?	Do not mix with sulfur products.
Thiolux (sulfur)	5-10 lb/acre	-	24	1	Not recommended within 2 weeks of an oil application nor if temperatures are expected to exceed 90 degrees within 3 days following

Table 7.4.4 Pesticides Labeled for Management of Powdery Mildew					
Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
					application.
Trilogy (neem oil)	1% solution	up to day	4	?	Maximum labeled use of 2 gal/acre/application
TriTek (mineral oil)	1-2 gal/100 gal water	up to day	4	?	Apply as needed.

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.
 PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.4.5 Gray Mold/Botrytis Fruit Rot (*Botrytis cinerea*)

Botrytis fruit rot usually begins as a small lesion at the blossom end or where a berry is touching another infected berry. The infected portion is firm and brown while the berry is still green, but it expands and softens as the fruit ripens. A powdery gray mass of spores covers infected berries if the weather remains humid and/or air circulation is poor.

IPM fact sheet on Gray Mold (Botrytis Fruit Rot) nysipm.cornell.edu/factsheets/berries/botrytis.pdf

Gray Mold (Botrytis Fruit Rot) Management Options	
Scouting/thresholds	None established.
Variety susceptibility	No known resistant varieties. Less severely impacted varieties are 'Earliglow', 'Jewel' and 'Clancy'. 'Allstar' and 'Sable' are very susceptible.
Cultural management	Disease control is greatly aided by managing weeds and by using other practices that promote good air circulation and rapid drying of the fruit such as regulating plant density. Use of protected production structures, such as low tunnels, reduces gray mold occurrence by limiting fruit wetness. Spring applications of nitrogen can dramatically increase the potential for infection. Prompt harvest of ripe fruit helps reduce disease development and spread. It may be beneficial to employ an hourly picker to remove only overripe and diseased fruit to prevent infection of clean fruit by other pickers. Overripe fruit should not be consumed. Cull piles should be buried or otherwise physically removed from fields during harvest.
Chemical treatment	Protection of blossoms is critical in gray mold management. Research in New York has consistently shown that excellent gray mold control can be obtained with just two fungicide sprays applied at early bloom and 10 days later. Continued protection of fruit prior to harvest may be necessary during prolonged periods of wet, foggy, or humid weather.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.4.5 Pesticides Labeled for Management of Gray Mold (Botrytis Fruit Rot)					
Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Actinovate AG (<i>Streptomyces Lydicus</i> WYEC 108)	3-12 oz/acre	0	until dry	2	For best results apply with a spreader/sticker prior to onset of disease.
Agricure (potassium bicarbonate)	2-5 lb/acre	0	1	?	
Cinnerate (cinnamon oil)	13-30 fl.oz./100 gal water	-	-	?	25(b) pesticide.
Cueva Fungicide Concentrate (copper octanoate)	0.5-2 gal/acre	up to day	4	1	
Double Nickel 55 (<i>Bacillus amyloliquefaciens</i> str D747)	0.25-3 lb/acre	0	4	2	
Double Nickel LC (<i>Bacillus amyloliquefaciens</i> str D747)	0.5-6 qt/acre	0	4	2	
JMS Stylet-Oil (paraffinic oil)	3 qt/100 gal water	0	4	2	A high volume of water is needed for thorough coverage. Many common pesticides are phytotoxic when applied with or close to oil sprays (e.g., sulfur). Check label for restrictions.
Milstop (potassium bicarbonate)	2-5 lb/acre	0	1	3	Do not mix with other pesticides or fertilizers. Not compatible with alkaline solutions.
Optiva (<i>Bacillus subtilis</i> str QST 713)	14-24 oz/acre	0	4	3	Repeat on 7-10 day intervals.
Oxdate 2.0 (hydrogen dioxide, peroxyacetic acid)	64 fl oz/100 gal water dip	0	until dry	?	Pre-plant dip.
Oxdate 2.0 (hydrogen dioxide, peroxyacetic acid)	32 fl oz – 1 gal/100 gal water	0	until dry	?	Foliar application, at-planting and for existing plantings. See label for additional instructions.
PERpose Plus (hydrogen peroxide)	1 fl oz/gal Initial/curative 0.25-33 fl oz/gal Weekly/preventative	-	until dry	3	For initial or curative use, apply higher rate for 1 to 3 consecutive days. Then follow with weekly/preventative treatment. For weekly or preventative treatments, apply lower rate every five to seven days. At first signs of disease, use curative rate then resume weekly preventative treatment.
Prestop (<i>Gliocladium catenulatum</i>)	3.5 oz/ 5 gal water	-	0	?	Apply only when no above-ground harvestable food commodities are present
PureSpray Green (white mineral)	0.75-1.5 gal/100	up to	4	?	Spray at no less than 400 PSI using

Table 7.4.5 Pesticides Labeled for Management of Gray Mold (Botrytis Fruit Rot)

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
oil)	gal water	day			ceramic nozzles.
Regalia (<i>Reynoutria sachalinensis</i>)	1-3 qt/acre	0	4	?	Initiate at first sign of disease then every 7-14 days.
Serenade ASO (<i>Bacillus subtilis</i>)	2-6 qt/acre	0	4	3	Begin application at or before flowering repeat every 7-10 days.
Serenade MAX (<i>Bacillus subtilis</i>)	1-3 lb/acre	0	4	3	See comment for Serenade ASO.
Serenade Opti (<i>Bacillus subtilis</i>)	14-20 oz/acre	0	4	3	See comment for Serenade ASO.
Serifel (<i>Bacillus amyloliquefaciens</i>)	4-16 oz/acre	0	4	?	
Trilogy (neem oil)	1% solution	up to day	4	?	Maximum labeled use of 2 gal/acre/application

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.

PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.4.6 Anthracnose (*Colletotrichum acutatum*)

One or more circular spots occur on the fruit. Spots originally are tan or light brown but become darker and sunken. Sunken spots are usually about one eighth to one quarter inch in diameter and may be covered with pink slimy spore masses during wet or very humid periods. The disease may occur on both green and ripe fruit, but is most common on ripe fruit following periods of warm, wet weather. In New York, anthracnose occurs only sporadically and is a more common problem on day-neutral varieties in the summer than it is on June-bearing varieties. However, the disease can be serious on June-bearing varieties if warm, wet weather conditions occur between fruit set and harvest.

Anthracnose Management Options

Scouting/thresholds	None established.
Variety susceptibility	No known resistant varieties. 'Jewel' shows little infection in field conditions.
Cultural management	Provide good air circulation by controlling weeds and reducing planting density. Use of protected production structures, such as low tunnels, reduces anthracnose occurrence by limiting fruit wetness. The anthracnose fungus is spread throughout a planting by splashing raindrops or sprinkler irrigation. Straw mulch may reduce the rate of disease spread relative to bare ground (less rain splash).
Chemical treatment	See table below.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.4.6 Pesticides Labeled for Management of Anthracnose

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Actinovate AG (<i>Streptomyces lydicus</i> WYEC 108)	3-12 oz/acre	0	until dry	2	For best results apply with a spreader/sticker prior to onset of disease. Re-apply as necessary.

Table 7.4.6 Pesticides Labeled for Management of Anthracnose

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Agricure (potassium bicarbonate)	2-5 lb/acre	0	1	?	
Cueva Fungicide Concentrate (copper octanoate)	0.5-2 gal/acre	up to day	4	1	
Double Nickel 55 (<i>Bacillus amyloliquefaciens</i> str D747)	0.25-3 lb/acre	0	4	2	
Double Nickel LC (<i>Bacillus amyloliquefaciens</i> str D747)	0.5-6 qt/acre	0	4	2	
Milstop (potassium bicarbonate)	2-5 lb/acre	0	1	?	Do not mix with other pesticides or fertilizers. Not compatible with alkaline solutions.
PERpose Plus (hydrogen peroxide)	1 fl oz/gal Initial/curative 0.25-33 fl oz/gal Weekly/pre-ventative	-	until dry	2	For initial or curative use, apply higher rate for 1 to 3 consecutive days. Then follow with weekly/preventative treatment. For weekly or preventative treatments, apply lower rate every five to seven days. At first signs of disease, use curative rate then resume weekly preventative treatment.
Regalia (<i>Reynoutria sachalinensis</i>)	1-3 qt/acre	0	4	?	Apply preventatively and repeat on a 7-10 day interval or as needed.
Serenade ASO (<i>Bacillus subtilis</i>)	2-6 qt/acre	0	4	2	Begin application at or before disease development then repeat every 7-10 days.
Serenade MAX (<i>Bacillus subtilis</i>)	1-3 lb/acre	0	4	2	Apply on a 7-10 schedule following disease onset.
Serenade Opti (<i>Bacillus subtilis</i>)	14-20 oz/acre	0	4	3	Begin application at or before flowering repeat every 7-10 days.
Serifel (<i>Bacillus amyloliquefaciens</i>)	4-16 oz/acre	0	4	?	
Trilogy (neem oil)	1% solution	Up to day	4	?	Maximum labeled use of 2 gal/acre/application

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.

PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.4.7 Leather Rot (*Phytophthora cactorum*)

Infected areas on immature fruit are brown, whereas those on maturing fruit appear bleached out. On all fruit, the infected areas are tough, leathery, and discolored on the inside as well as the outside of the fruit. Diseased fruits have a pungent smell and bitter taste. Leather rot is most severe during periods of abundant warm rains during the fruiting period and in flooded soils. The cultural practices listed in the table below are the most effective control procedures.

IPM fact sheet Leather Rot nysipm.cornell.edu/factsheets/berries/leather_rot.pdf

Leather Rot Management Options	
Scouting/thresholds	None established.
Variety susceptibility	No known resistant varieties.
Cultural management	<p>Plant only on a well-drained site or provide supplemental drainage. Growing strawberries on raised beds will also reduce disease severity.</p> <p>Minimize soil flooding through site selection; by avoiding planting in ruts; and by preventing or reducing soil compaction.</p> <p>Provide an extra layer of straw mulch between rows throughout the fruiting season. The mulch provides a physical barrier between the soilborne pathogen and the susceptible fruit.</p>
Chemical treatment	See below.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.4.7 Pesticides Labeled for Management of Leather Rot

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Actinovate AG (<i>Streptomyces Lydicus</i> WYEC 108)	3-12 oz/acre soil drench	0	until dry	?	Since Actinovate AG contains live spores of a microbe, best results will be obtained if used prior to disease onset.
BIO-TAM (<i>Trichoderma asperellum</i> , <i>Trichoderma gamsii</i>)	0.25-2 lb/gal water dip	-	1	?	Bare root dip
BIO-TAM (<i>Trichoderma asperellum</i> , <i>Trichoderma gamsii</i>)	1.5-3 oz/1000 row feet in-furrow treatment	-	1	?	
BIO-TAM (<i>Trichoderma asperellum</i> , <i>Trichoderma gamsii</i>)	2.5-3 lb/acre band	-	1	?	
Bio-Tam 2.0 (<i>Trichoderma asperellum</i> , <i>Trichoderma gamsii</i>)	2.5-5 lb/acre	-	1	?	
Double Nickel 55 (<i>Bacillus amyloliquefaciens</i> str D747)	0.125-1 lb/acre Soil treatment	0	4	?	
Double Nickel LC (<i>Bacillus amyloliquefaciens</i> str D747)	0.5-4.5 pts/acre Soil treatment	0	4	?	
PERpose Plus (hydrogen peroxide)	<p>1 fl oz/gal Initial/curative</p> <p>0.25-33 fl oz/gal Weekly/preventative</p>	-	until dry	?	<p>For initial or curative use, apply higher rate for 1 to 3 consecutive days. Then follow with weekly/preventative treatment.</p> <p>For weekly or preventative treatments, apply lower rate every five to seven days. At first signs of disease, use curative rate then resume weekly preventative</p>

Table 7.4.7 Pesticides Labeled for Management of Leather Rot

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
					treatment.
Prestop (<i>Gliocladium catenulatum</i>)	1.4-3.5 oz/ 2.5 gal water soil drench	-	0	?	Treat only the growth substrate when above-ground harvestable food commodities are present.
RootShield PLUS+ Granules (<i>Trichoderma harzianum</i> , <i>Trichoderma virens</i>)	2.5-6 lb/half acre in-furrow treatment	-	0	?	
RootShield PLUS+ WP (<i>Trichoderma harzianum</i> , <i>Trichoderma virens</i>)	0.25-5 lb/ 20 gal water dip	0	4	?	Do not apply when above-ground harvestable food commodities are present.
RootShield PLUS+ WP (<i>Trichoderma harzianum</i> , <i>Trichoderma virens</i>)	16-32 oz/acre in-furrow treatment	0	4	?	Do not apply when above-ground harvestable food commodities are present.
TerraClean 5.0 (hydrogen dioxide, peroxyacetic acid)	25 fl oz/ 200 gal water/1,000 sq ft soil treated soil drench	up to day	0	?	See label for rate information for specific soil treatments.
Zonix (<i>Rhamnolipid Biosurfactant</i>)	0.5-0.8 fl.oz./ gal water	-	4	?	Prepare enough solution based on plant density and soil conditions to insure thorough coverage.

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.

PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.4.8 Red Stele (*Phytophthora fragariae*)

Red stele is caused by a soilborne aquatic pathogen (*Phytophthora*) that may persist in the soil for many years even when strawberries are no longer grown. Symptoms of infection often appear just before harvest. Diseased plants appear stunted and off-color, and will often wilt and collapse if the weather becomes warm and dry. Because these same symptoms may be caused by other factors that destroy roots (such as root-feeding insects), the diagnosis depends on an examination of the plant's root system. In a diseased plant, the roots have a "rat-tail" appearance caused by loss of the fine branched feeder roots from the main fleshy roots. The main fleshy roots are rotted from the tips back toward the crown. Cutting or scraping away the white outer portion (epidermis and cortex) just above the rotten areas in early infections sometimes reveals a reddish root core (stela). Infected plants usually appear in groups and are frequently found in the lowest or wettest parts of a field.

IPM fact sheet Red Stele nysipm.cornell.edu/factsheets/berries/red_stele.pdf

Red Stele Management Options	
Scouting/thresholds	None established.
Variety susceptibility	Resistant varieties include 'Earliglow', 'Northeast', 'Mohawk', 'Redchief', 'Guardian', 'Allstar', 'Tribute', 'Tristar', 'Surecrop', and 'Sparkle'. However, these varieties are not resistant to all races of the red stele pathogen (<i>P. fragariae</i>), and as such, the disease could still develop if a race to which they are not resistant is present. 'Jewel' and 'Honeye' are especially susceptible.
Cultural management	Because the red stele fungus is particularly active in extremely wet soil, plant only on a well-drained site or provide supplemental drainage. Growing strawberries on raised beds will also reduce disease severity.

Red Stele Management Options	
Chemical treatment	The red stele fungus is not present in every field, thus treatments should be confined to fields and areas within fields where the disease has occurred previously or is suspected.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.4.8 Pesticides Labeled for Management of Red Stele					
Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Actinovate AG (<i>Streptomyces Lydicus</i> WYEC 108)	3-12 oz/acre soil drench	0	until dry	?	Since Actinovate AG contains live spores of a microbe, best results will be obtained if used prior to disease onset.
BIO-TAM (<i>Trichoderma asperellum</i> , <i>Trichoderma gamsii</i>)	0.25-2 lb/ gal water dip	-	1	?	Bare root dip.
BIO-TAM (<i>Trichoderma asperellum</i> , <i>Trichoderma gamsii</i>)	1.5-3 oz/1000 row feet in-furrow	-	1	?	
BIO-TAM (<i>Trichoderma asperellum</i> , <i>Trichoderma gamsii</i>)	2.5-3 lb/acre band	-	1	?	
Bio-Tam 2.0 (<i>Trichoderma asperellum</i> , <i>Trichoderma gamsii</i>)	2.5-3 lb/acre	-	1	?	
Double Nickel 55 (<i>Bacillus amyloliquefaciens</i> str D747)	0.125-1 lb/acre Soil treatment	0	4	?	
Double Nickel LC (<i>Bacillus amyloliquefaciens</i> str D747)	0.5-4.5 pts/acre Soil treatment	0	4	?	
PERpose Plus (hydrogen peroxide)	1 fl oz/gal Initial/curative 0.25-33 fl oz/gal Weekly/preventative	-	until dry	?	For initial or curative use, apply higher rate for 1 to 3 consecutive days. Then follow with weekly/preventative treatment. For weekly or preventative treatments, apply lower rate every five to seven days. At first signs of disease, use curative rate then resume weekly preventative treatment.
Prestop (<i>Gliocladium catenulatum</i>)	1.4-3.5 oz/ 2.5 gal water soil drench	-	0	?	Treat only the growth substrate when above-ground harvestable food commodities are present.
Regalia (<i>Reynoutria sachalinensis</i>)	1-4 qt/100 gal water dip	0	4	?	Pre-plant dip.
RootShield PLUS+ Granules (<i>Trichoderma harzianum</i> , <i>Trichoderma virens</i>)	2.5-6 lb/half acre in-furrow treatment	-	0	?	
RootShield PLUS+ WP (<i>Trichoderma harzianum</i> ,	0.25-5 lb/ 20 gal water dip	0	4	?	Do not apply when above-ground harvestable food commodities are

Table 7.4.8 Pesticides Labeled for Management of Red Stele

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
<i>Trichoderma virens</i>)					present.
RootShield PLUS+ WP (<i>Trichoderma harzianum</i> , <i>Trichoderma virens</i>)	16-32 oz/acre in-furrow treatment	0	4	?	See previous RootShield Plus+ WP comment.
TerraClean 5.0 (hydrogen dioxide, peroxyacetic acid)	25 fl.oz./ 1,000sq ft soil treated soil drench	up to day	0	?	See label for rate information for specific soil treatments.
Zonix (Rhamnolipid Biosurfactant)	0.5-0.8 fl.oz./ gal water	-	4	?	Prepare enough solution based on plant density and soil conditions to insure thorough coverage

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.

PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.4.9 Black Root Rot

Black root rot constitutes a complex set of symptoms caused by one or more of the following organisms: nematodes, root rot fungi (*Pythium spp.*, *Rhizoctonia spp.*). Black root rot is most commonly observed in older plantings or on heavy compacted soils. Over time, plant vigor and productivity declines. Feeder rootlets die, and fleshy structural roots deteriorate and become blackened. The blackening starts as patches along the length of the root, rather than from the tip back. This disease is often associated with fields having a long history of strawberry production. Because no single cause of black root rot has been defined, there is no single control.

Black Root Rot Management Options

Scouting/thresholds	None established.
Variety susceptibility	No known resistant varieties. Particularly susceptible varieties are 'Honeoye' and 'Jewel'. These varieties should be avoided in fields without a adequate rotation.
Cultural management	Fields with high nematode populations may be more prone to black root rot development. Check nematode populations prior to planting. If high, consider incorporating a cover crop with biofumigant properties. See Section 4, Cover Crops. Cultural practices that reduce soil compaction, improve aeration, and promote good drainage are beneficial for reducing disease. Rotating a field out of strawberries for at least 2 - 3 years is strongly recommended. Measures to control red stele will also help alleviate black root rot. Cover crops such as brown mustard and indiangrass and incorporation of compost can also provide disease suppression.
Chemical treatment	Chemical treatment with a nematicide is not suggested for black root rot. See below for fungicides.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.4.9 Pesticides Labeled for Management of Black Root Rot

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Actinovate AG (<i>Streptomyces Lydicus</i> WYEC 108)	3-12 oz/acre soil drench	0	until dry	?	Since Actinovate AG contains live spores of a microbe, best results will be obtained if used prior to disease onset.
BIO-TAM (<i>Trichoderma asperellum</i> , <i>Trichoderma gamsii</i>)	0.25-2 lb/ gal water dip	-	1	?	Bare root dip.
BIO-TAM (<i>Trichoderma asperellum</i> , <i>Trichoderma gamsii</i>)	1.5-3 oz/1000 row feet in-furrow treatment	-	1	?	
BIO-TAM (<i>Trichoderma asperellum</i> , <i>Trichoderma gamsii</i>)	2.5-3 lb/acre band	-	1	?	
Bio-Tam 2.0 (<i>Trichoderma asperellum</i> , <i>Trichoderma gamsii</i>)	2.5-3 lb/acre	-	1	?	
Double Nickel 55 <i>Bacillus amyloliquefaciens</i> str D747)	0.125-1 lb/acre Soil treatment	0	4	?	
Double Nickel LC (<i>Bacillus amyloliquefaciens</i> str D747)	0.5-4.5 pts/acre Soil treatment	0	4	?	
PERpose Plus (hydrogen peroxide)	1 fl oz/gal Initial/curative 0.25-33fl oz/gal Weekly/pre-ventative	-	until dry	?	For initial or curative use, apply higher rate for 1 to 3 consecutive days. Then follow with weekly/preventative treatment. For weekly or preventative treatments, apply lower rate every five to seven days. At first signs of disease, use curative rate then resume weekly preventative treatment.
Prestop (<i>Gliocladium catenulatum</i>)	1.4-3.5 oz/ 2.5 gal water soil drench	-	0	?	Treat only the growth substrate when above-ground harvestable food commodities are present.
Regalia (<i>Reynoutria sachalinensis</i>)	1-3 qt/100 gal water soil drench	0	4	?	
Regalia (<i>Reynoutria sachalinensis</i>)	1-4 qt/100 gal water dip	0	4	?	Pre-plant dip.
RootShield Granules (<i>Trichoderma harzianum</i>)	2.5-6 lb/half acre in-furrow	-	0	?	
RootShield PLUS+ Granules (<i>Trichoderma harzianum</i> , <i>Trichoderma virens</i>)	2.5-6 lb/half acre in-furrow	-	0	?	
RootShield PLUS+ WP (<i>Trichoderma</i>	0.25-5 lb/ 20 gal	0	4	?	

Table 7.4.9 Pesticides Labeled for Management of Black Root Rot

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
<i>harzianum</i> , <i>Trichoderma virens</i>)	water dip				
RootShield PLUS+ WP (<i>Trichoderma harzianum</i> , <i>Trichoderma virens</i>)	16-32 oz/acre in-furrow treatment	0	4	?	
RootShield WP (<i>Trichoderma harzianum</i>)	3-5 oz/100 gal water soil drench	-	until dry	?	Greenhouse/nursery drench. No PHI specified on the label for any of the product rates.
RootShield WP (<i>Trichoderma harzianum</i>)	0.5-2.5 lb/ 5 gal dip	-	until dry	?	Bare root dip. No PHI specified on the label for any of the product rates.
RootShield WP (<i>Trichoderma harzianum</i>)	16-32 oz/acre in-furrow treatment	-	until dry	?	In-furrow or transplant starter solution. No PHI specified on the label for any of the product rates.
Soilgard (<i>Gliocladium virens</i>)	2-10 lb/acre	-	0	?	
TerraClean 5.0 (hydrogen dioxide, peroxyacetic acid)	25 fl oz/ 200 gal water/1,000 sq ft soil treated soil drench	Up to day	0	?	See label for rate information for specific soil treatments.

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.

PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.5 OTHER DISEASES OF NOTE

7.5.1 Angular Leaf Spot (*Xanthomonas fragariae*)

Tiny water-soaked lesions appear first on lower leaf surfaces. These enlarge to form angular spots usually bordered by small veins. When held up to the light spots appear translucent, but are dark green under reflected light. Spots may ooze bacteria under moist conditions, which dry to form a whitish scaly skin. Lesions eventually become visible on upper leaf surfaces as irregular reddish brown spots. Calyxes may also become infected. The disease is favored by daytime temperatures around 68°F, low to near freezing night temperatures, and precipitation events such as rain, overhead irrigation or heavy dews.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.5.1 Pesticides Labeled for Management of Angular Leaf Spot

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Badge X2 (copper hydroxide, copper oxychloride)	1-2.5 lb/acre	0	48	?	
Champ WG (copper hydroxide)	2-3 lb/acre	-	48	?	May cause crop injury under some conditions.
CS 2005 (copper sulfate pentahydrate)	19.2-25.6 oz/acre	0	48	?	
Cueva Fungicide Concentrate (copper octanoate)	0.5-2 gal/acre	up to day	4	?	
Double Nickel 55 (<i>Bacillus</i>	0.25-3 lb/acre	0	4	?	

Table 7.5.1 Pesticides Labeled for Management of Angular Leaf Spot					
<i>amyloliquefaciens</i> str D747)					
Double Nickel LC (<i>Bacillus amyloliquefaciens</i> str D747)	0.5-6 qt/acre	0	4	?	
Oxidate 2.0 (hydrogen dioxide, peroxyacetic acid)	32 fl oz – 1 gal/100 gal water	0	until dry	?	
PERpose Plus (hydrogen peroxide)	1 fl oz/gal Initial/curative 0.25-33fl oz/gal Weekly/preventative	-	until dry	3	For initial or curative use, apply higher rate for 1 to 3 consecutive days. Then follow with weekly/preventative treatment. For weekly or preventative treatments, apply lower rate every five to seven days. At first signs of disease, use curative rate then resume weekly preventative treatment.
Trilogy (neem oil)	1% solution	up to day	4	?	Maximum labeled use of 2 gal/acre/application

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.

PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.5.2 Verticillium Wilt (*Verticillium albo-atrum*)

Plants are affected most severely during their first year of growth. Outer leaves turn brown and eventually collapse, but inner leaves remain green until the plant dies. This symptom distinguishes Verticillium wilt from other root and crown disorders. Affected plants may occur uniformly, but more typically, they appear scattered throughout a field. In problem areas or after the last crop of tomatoes, potatoes, or eggplant, plant only varieties resistant to Verticillium wilt for at least 3 years. Resistant varieties include 'Earliglow', 'Guardian', 'Allstar', 'Tribute', and 'Tristar'. Many weeds are hosts of the Verticillium fungus, particularly nightshade, groundcherry, redroot pigweed, lambsquarters, and horsenettle. These weeds should be strictly controlled in current and future planting sites to keep *Verticillium* inoculum low.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.5.2 Pesticides Labeled for Management of Verticillium Wilt					
Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Actinovate AG (<i>Streptomyces lydicus</i> WYEC 108)	3-12 oz/acre soil drench	0	until dry	?	Since Actinovate AG contains live spores of a microbe, best results will be obtained if used prior to disease onset.
BIO-TAM (<i>Trichoderma asperellum</i> , <i>Trichoderma gamsii</i>)	0.25-2 lb/ gal water dip	-	1	?	Bare root dip.
BIO-TAM (<i>Trichoderma asperellum</i> , <i>Trichoderma gamsii</i>)	1.5-3 oz/1000 row feet in-furrow	-	1	?	
BIO-TAM (<i>Trichoderma asperellum</i> , <i>Trichoderma gamsii</i>)	2.5-3 lb/acre band	-	1	?	

Table 7.5.2 Pesticides Labeled for Management of Verticillium Wilt					
Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Bio-Tam 2.0 (<i>Trichoderma asperellum</i> , <i>Trichoderma gamsii</i>)	2.5-3 lb/acre	-	1	?	
Double Nickel 55 (<i>Bacillus amyloliquefaciens</i> str D747)	0.125-1 lb/acre soil treatment	0	4	?	
Double Nickel LC (<i>Bacillus amyloliquefaciens</i> str D747)	0.5-4.5 pts/acre soil treatment	0	4	?	
PERpose Plus (hydrogen peroxide)	1 fl oz/gal Initial/curative	-	until dry	?	For initial or curative use, apply higher rate for 1 to 3 consecutive days. Then follow with weekly/preventative treatment.
PERpose Plus (hydrogen peroxide)	0.25-33fl oz/gal Weekly/preventative	-	until dry	?	For weekly or preventative treatments, apply lower rate every five to seven days. At first signs of disease, use curative rate then resume weekly preventative treatment.
Prestop (<i>Gliocladium catenulatum</i>)	1.4-3.5 oz/ 2.5 gal water soil drench	-	0	?	Treat only the growth substrate when above-ground harvestable food commodities are present.
Regalia (<i>Reynoutria sachalinensis</i>)	1-4 qt/100 gal water dip	0	4	?	Pre-plant dip.
Serenade Soil (<i>Bacillus subtilis</i>)	2-6 qt/acre soil treatment	0	4	?	Soil drench or in-furrow.
TerraClean 5.0 (hydrogen dioxide, peroxyacetic acid)	25 fl oz/ 200 gal water/1,000sq ft soil treated soil drench	up to day	0	?	See label for rate information for specific soil treatments.

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.

PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.6 INSECTS AND MITES OF PRIMARY CONCERN

The insects and mites that are considered major pests in strawberries can vary in occurrence both from year to year and from site to site. For these reasons it is important to be familiar with the life cycle of the pest to assist in developing a scouting program that will ensure a pest problem can be discovered and dealt with before it becomes an outbreak. Alternatively, it is important to know when a potential pest is not causing significant economic damage so that unnecessary controls can be avoided. Applying an organically approved broad-spectrum insecticide such as PyGanic EC (a pyrethrum) when not necessary, for example, is not only a waste of money but also has the potential to disrupt biological control by beneficial organisms. This illustrates the need to take potential biological control agents (predators, parasitoids, parasites, microbes) into account when making management decisions. Following are descriptions of the most commonly found insect pests in strawberry plantings.

7.6.1 Root Weevil (*various species*)

Different species, but most commonly the strawberry root weevil, the black vine weevil, and the rough strawberry root weevil. These pests attack the roots or crowns of plants while in the grub stage. All have a one-year life cycle, although some are known to live two seasons. Adults emerge about late June. Adults are active at night and feed on leaves, causing

characteristic semi-circle leaf chewing on leaf edges. This feeding damage is not considered of economic importance. Adults can not fly. Beds with heavy infestations show distinct patches or spots that appear stunted and have substantially reduced yields. The roots of injured plants are badly eaten away, and continued infestation may destroy infested plants.

IPM fact sheet Root Weevil nysipm.cornell.edu/factsheets/berries/root_weevils.pdf

Root Weevil Management Options	
Scouting/thresholds	None established.
Variety susceptibility	None adapted to the Northeastern region.
Cultural management	Rotate out of strawberries for at least 1 year to reduce root weevil density. A barrier (plastic fence) can prevent walking adults (they can not fly) from moving from an infested field to a new field to be planted. See Exclusion Barriers for Management of Black Vine Weevil for details.
Biological control	Two species of <i>Heterorhabditis</i> , insect parasitic nematodes, <i>H. bacteriophora</i> and <i>H. marelatus</i> , can provide control of larvae. Release nematodes either in spring when soils warm (>50 F) or in late summer - early fall. Provide sufficient water to move nematodes into the root zone. Applications may need to be repeated annually.
Chemical treatment	Available chemical treatments target adults. See below.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.6.1 Pesticides Labeled for Management of Root Weevil					
Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Aza-Direct (azadirachtin)	1-2 pts/acre	0	4	?	Foliar spray or soil drench.
AzaGuard (azadirachtin)	10-16 fl oz/acre	0	4	?	Apply with OMRI approved spray oil. Foliar spray or soil drench.
AzaMax (azadirachtin)	1.33 fl oz/ 1000 ft ²	0	4	?	Foliar spray or soil drench.
AzaSol (azadirachtin)	6 oz/acre	-	4	?	
Azatrol-EC (azadirachtin)	0.29-0.96 fl oz/ 1000 ft ²	0	4	?	
Azera (azadirachtin, pyrethrins)	1-3.5 pts/acre	-	12	?	Foliar spray or soil drench.
BioLink (garlic juice)	0.5-2 qt/acre	-	-	?	25(b) pesticide.
BioLink Insect & Bird Repellent (garlic juice)	0.5-4 qt/acre	-	-	?	25(b) pesticide.
Ecozin Plus 1.2% ME (azadirachtin)	15-30 oz/acre	0	4	?	Foliar spray or soil drench.
Envirepel 20 (garlic juice)	10-32 oz/acre	-	-	?	25(b) pesticide. Repellent.
Garlic Barrier AG (garlic juice)	See comments	-	-	?	25(b) pesticide. See label for specific information.
Molt-X (azadirachtin)	10 oz/acre	0	4	?	Foliar spray or soil drench.

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.

PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.6.2 STRAWBERRY SAP BEETLE (*Stelidota geminata*) & PICNIC BEETLE (*Glischrochilus fasciatus*)

Sap beetle adults make cavities in ripe and overripe fruit as well as spread spores of decay organisms. The larvae also feed on ripe and overripe fruit and are a source of contamination in harvested fruit. Until a few years ago, sap beetles were uncommon in strawberries. Now, sap beetles are occasionally found in high numbers in later ripening strawberry plantings throughout the state. Two species feed on strawberry fruits: the common picnic beetle, one quarter inch long with four yellow spots on the back, and the smaller, brown strawberry sap beetle without distinctive markings. Strawberry sap beetle is the more serious pest because it does not limit its activity to over-ripe fruit. Beetles overwinter at the edge of woodlots and possibly under other perennial fruit crops, such as brambles and blueberries, but they do not appear to overwinter in strawberry fields. As strawberries ripen, beetles move into the field and begin feeding and laying eggs. Fruit touching the ground or straw mulch appears particularly vulnerable. Adult strawberry sap beetle are very secretive and scamper away when disturbed, therefore they are sometimes hard to spot even when feeding damage is evident.

IPM fact sheet Sap Beetle nysipm.cornell.edu/factsheets/berries/ssb.pdf

Sap Beetle Management Options	
Scouting/thresholds	None established.
Variety susceptibility	No known resistant varieties, although cultivars that tend to hold fruit off the ground may be less vulnerable to adult feeding and larval contamination.
Cultural management	Keep the field free of ripe and over-ripe fruit. Good sanitation in berry crops and other fruit crops on the farm will help reduce food resources.
Chemical treatment	Generally insecticide sprays are not very effective, partly because it is difficult to get material where the insects are active (underside of fruit touching the ground). Therefore, good coverage is important and even then, it's unclear if level of control justifies the cost.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](https://www.pims.ny.gov/)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.6.2 Pesticides Labeled for Management of Sap Beetle					
Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Aza-Direct (azadirachtin)	1-2 pts/acre	0	4	3	Foliar spray or soil drench.
AzaGuard (azadirachtin)	8-16 fl.oz./acre	0	4	3	Apply with OMRI approved spray oil. Foliar spray or soil drench.
AzaMax (azadirachtin)	1.33 fl oz/1000 sq ft	0	4	3	Foliar spray or soil drench.
AzaSol (azadirachtin)	6 oz/acre	-	4	3	
Azatrol-EC (azadirachtin)	0.29-0.96 fl oz/1000 sq ft	0	4	3	Foliar spray or soil drench.
Azera (azadirachtin, pyrethrins)	1-3.5 pts/acre	-	12	?	Foliar spray or soil drench.
BioLink (garlic juice)	0.5-2 qt/acre		-	?	25(b) pesticide.
BioLink Insect & Bird Repellent (garlic juice)	0.5-4 qt/acre	-	-	?	25(b) pesticide.
Ecozin Plus 1.2% ME (azadirachtin)	15-30 oz/acre	0	4	3	Foliar spray or soil drench.
Envirepel 20 (garlic juice)	10-32 oz/acre	-	-	?	25(b) pesticide. Repellent.

Table 7.6.2 Pesticides Labeled for Management of Sap Beetle

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Garlic Barrier AG (garlic juice)	See comments.	-	-	?	25(b) pesticide. See label for specific information.
Molt-X (azadirachtin)	8 oz/acre	0	4	3	Foliar spray or soil drench.
PyGanic EC 1.4 II (pyrethrins)	16-64 fl oz/acre	until dry	12	?	Short residual activity may require multiple applications. Caution: do not use when bees are active in the planting.
PyGanic EC 5.0 II (pyrethrins)	4.5-17 fl oz/acre	until dry	12	?	See comment for PyGanic EC 1.4 II

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.

PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.6.3 TARNISHED PLANT BUG (*Lygus lineolaris*)

This pest causes “cat faced” or “button” berries. It damages the fruit by feeding on the developing fruit. The fruit tissue in the immediate area of damaged seeds stops developing. Little information is available on cultivar differences in susceptibility to tarnished plant bug, but early maturity is correlated with freedom from injury; later cultivars may suffer more damage. Also, highly productive cultivars appear to tolerate feeding damage better than less productive ones. Tarnished plant bug feeds on many crop and non-crop plants as they flower and fruit. Hence, weedy fields can promote higher populations. Also, populations increase during the season (there are 2 to 3 generations each year). As such, day-neutral strawberries in late summer often experience high levels of tarnished plant bug damage.

IPM fact sheet Tarnished Plant Bug nysipm.cornell.edu/factsheets/berries/tpb.pdf

Tarnished Plant Bug Management Options

Scouting/thresholds	Anytime from just before the blossoms open until harvest, check for tarnished plant bug nymphs by striking the plant over a flat, low-sided, light-colored dish. Suggested action threshold: 0.5 nymphs per cluster, or 4 out of 15 clusters with 1 or more nymphs.
Variety susceptibility	‘Honeoye’ and other highly productive cultivars appear less susceptible to feeding injury. Early-flowering cultivars may be less susceptible to injury also. Day-neutral varieties are particularly vulnerable later in the season.
Cultural management	Row covers accelerate plant development and help avoid injury. Pressure is often highest in weedy fields or in fields bordered by woody shrubs.
Chemical treatment	See below.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide’s effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](https://pims.ny.gov/)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.6.3 Pesticides Labeled for Management of Tarnished Plant Bug

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Aza-Direct (azadirachtin)	1-2 pts/acre	0	4	1	
AzaGuard (azadirachtin)	10-16 fl oz/acre	0	4	1	Apply with OMRI approved spray oil.

Table 7.6.3 Pesticides Labeled for Management of Tarnished Plant Bug

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
AzaMax (azadirachtin)	1.33 fl oz/1000 sq ft	0	4	1	
AzaSol (azadirachtin)	6 oz/acre	-	4	1	For nymph treatment.
Azatrol-EC (azadirachtin)	0.24-0.96 fl oz/1000 sq ft	0	4	1	
Azera (azadirachtin, pyrethrins)	1-3.5 pts/acre	-	12	1	
DES-X (insecticidal soap)	2% solution sprayed at 75-200 gallons/acre	1/2	12	?	
Ecotec (rosemary oil, peppermint oil)	1-4 pt/acre	-	-	?	25(b) pesticide.
Ecozin Plus 1.2% ME (azadirachtin)	15-30 oz/acre	0	4	1	
Envirepel 20 (garlic juice)	10-32 oz/acre	-	-	?	25(b) pesticide. Repellant.
Garlic Barrier AG (garlic juice)	See comments	-	-	?	25(b) pesticide. See label for specific information.
Grandevo (<i>Chromobacterium subtsugae</i> str PRAA4-1)	2-3 lb/acre	0	4	?	
Molt-X (azadirachtin)	10 oz/acre	0	4	1	
PFR-97 20% WDG (<i>Isaria fumosorosea</i> <i>Apopka</i> str 97)	1-2 lb/acre	-	4	?	
PyGanic EC 1.4 II (pyrethrins)	16-64 fl oz/acre	until dry	12	?	Short residual activity may require multiple applications. Caution: do not use when bees are active in the planting.
PyGanic EC 5.0 II (pyrethrins)	4.5-17 fl oz/acre	until dry	12	?	See comment for Pyganic 1.4 II

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.

PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.6.4 TWO-SPOTTED SPIDER MITE (*Tetranychus urticae*)

In early spring, mites begin feeding on the undersides of new leaves, sometimes resulting in small yellow spots on the upper leaf surfaces. These symptoms do not occur in all cases. Brownish dry areas on the lower leaf surfaces are more characteristic of damage. Later, the entire lower leaf may become dry and brown, giving it a bronzed appearance. Heavily infested plants look dry and stunted, and their sparse new growth is yellowish and distorted. Damage is first seen and is most prevalent in dry areas of a field. Mild growing areas in New York (Hudson Valley and Long Island) experience problems with mites most frequently.

Two-spotted Spider Mite Management Options	
Scouting/thresholds	Five mites/leaf or 15 out of 60 mature (fully expanded) leaflets infested with 1 or more mites. Regular leaf monitoring is necessary for assessing population growth.
Variety susceptibility	No known resistant varieties.

Two-spotted Spider Mite Management Options	
Cultural management	<p>Ensure plots are not over fertilized.</p> <p>Provide adequate irrigation. Cool, moist conditions are unfavorable to mites.</p> <p>Do not use other insecticides that kill predatory mites.</p> <p>Mow and incorporate leaves at renovation.</p>
Biological Control	<p>Species of predatory mites can be purchased from biological control supply companies and released into strawberry fields to provide some control of spider mites.</p> <p>Effectiveness has not been carefully assessed under NY conditions. Note that predatory mites should be released before significant feeding damage is observed. <i>Neoseiulus californicus</i> and <i>Amblyseius fallacis</i> are two predatory mite species used in strawberries for biological control of two-spotted spider mite.</p>
Chemical treatment	<p>Chemical control of spider mites is often not completely effective because of their high mobility, tendency to reside on the underside of leaves where it is difficult to reach with miticides, high reproductive rate, and resistance to some pesticides. Good coverage of the plants, particularly the undersides of the leaves, is critical for adequate protection. Use adequate water (200 - 300 gal/A) for maximum effectiveness of the miticide. Repeat at 7- to 10-day intervals as necessary unless otherwise noted on label.</p> <p>Soap sprays may provide some control but excellent coverage is essential, especially on lower leaf surfaces.</p>

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.6.4 Pesticides Labeled for Management of Two-spotted Spider Mite					
Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Acoidal (sulfur)	5-10 lb/acre	-	24	1	Do not use on sulfur sensitive varieties.
Aza-Direct (azadirachtin)	1-2 pts/acre	0	4	1	
AzaGuard (azadirachtin)	10-16 fl oz/acre	0	4	1	Apply with OMRI approved spray oil.
AzaMax (azadirachtin)	1.33 fl oz/1000 ft ²	0	4	1	
Azatrol-EC (azadirachtin)	0.24-0.96 fl oz/1000 ft ²	0	4	1	
Azera (azadirachtin, pyrethrins)	1-3.5 pts/acre	-	12	1	
BioLink (garlic juice)	0.5-2 qt/acre	-	-	?	25(b) pesticide.
BioLink Insect & Bird Repellent (garlic juice)	0.5-4 qt/acre	-	-	?	25(b) pesticide.
Cinnerate (cinnamon oil)	13-30 fl oz/100 gal water	-	-	?	25(b) pesticide.
Defend DF (sulfur)	5-10 lb/acre	-	24	1	Do not use on sulfur sensitive varieties.
DES-X (insecticidal soap)	2% solution sprayed at	1/2	12	?	

Table 7.6.4 Pesticides Labeled for Management of Two-spotted Spider Mite

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
	75-200 gallons/acre				
Ecotec (rosemary oil, peppermint oil)	1-4 pt/acre	-	-	3	25(b) pesticide. Essential oils effective against spider mites in 0/1 trial.
Envirepel 20 (garlic juice)	10-32 oz/acre	-	-	?	25(b) pesticide. Repellant.
Garlic Barrier AG (garlic juice)	See comments	-	-	?	25(b) pesticide. See label for specific information.
GC-Mite (garlic oil, clove oil, cottonseed oil)	1 gal/100 gal water	-	-	1	25(b) pesticide. Conduct compatibility test prior to application.
Glacial Spray Fluid (mineral oil)	0.75 gal/ 100 gal	up to day	4	1	See label for specific application volumes and equipment.
Grandevo (<i>Chromobacterium subtsugae</i> str PRAA4-1)	2-3 lb/acre	0	4	?	
GrasRoots (cinnamon oil)	1 part GrasRoots: 9 parts water	-	-	?	25(b) pesticide.
JMS Stylet-Oil (paraffinic oil)	3 qt/100 gal water	0	4	1	Apply for optimum coverage of leaf surfaces. Use high pressure, small droplet size, and adequate gallonage to ensure good coverage. Can cause phytotoxicity if applied too close to a sulfur application.
Micro Sulf (sulfur)	5-10 lb/acre	-	24	?	Some varieties may be sensitive to sulfur.
Microthiol Disperss (sulfur)	5-10 lb/acre	-	24	?	Not recommended within 2 weeks of an oil application nor if temperatures are expected to exceed 90 degrees within 3 days following the application.
M-Pede (insecticidal soap)	1-2% vol/vol	0	12	1	Works by contact. Good coverage is important.
Nuke Em (citric acid)	1 fl oz/ 31 fl oz water. Normal strength	-	-	?	25(b) pesticide. Use the normal strength mix first. See label for stronger dilutions if needed.
Oleotrol-I Bio-Insecticide Concentrate (soybean oil)	43-45 fl.oz./100 gal water	-	-	?	25(b) pesticide.
Omni Supreme Spray (mineral oil)	1-2% vol/vol	-	12	1	See label for specific precautions. Applied at 60 gallons of finished spray per acre when using air-assisted, low-volume ground application equipment or 200 gallons of water per acre with standard ground spray equipment.

Table 7.6.4 Pesticides Labeled for Management of Two-spotted Spider Mite

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Organocide (sesame oil)	1-2 gal/100 gal water	-	-	1	25(b) pesticide.
PFR-97 20% WDG (Isaria fumosorosea Apopka str. 97)	1-2 lb/acre	-	4	?	
PureSpray Green (white mineral oil)	0.75-1.5 gal/100 gal water	up to day	4	1	Spray at no less than 400 PSI using ceramic nozzles.
PyGanic EC 1.4 II (pyrethrins)	16-64 fl.oz./acre	until dry	12	?	Short residual activity may require multiple applications. Caution: do not use when bees are active in the planting.
PyGanic EC 5.0 II (pyrethrins)	4.5-17 fl.oz./acre	until dry	12	?	See comment for PyGanic EC 1.4 II
Sil-Matrix (potassium silicate)	0.5-1% solution.	0	4	?	
SuffOil-X (aliphatic petroleum solvent)	1-2 gal/100 gal water	up to day	4	1	Do not mix with sulfur products.
Trilogy (neem oil)	1-2% solution	up to day	4	?	Maximum labeled use of 2 gal/acre/application
TriTek (mineral oil)	1-2 gal/100 gal water	up to day	4	1	Apply as needed.

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.

PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.6.5 SPOTTED WING DROSOPHILA (*Drosophila suzukii*)

Spotted Wing Drosophila (SWD) is an invasive vinegar or fruit fly that was first detected in NY in 2011 and spread across NY in 2012. June-bearing strawberries and day neutral strawberries fruiting early in the growing season have thus far escaped injury from this pest in NY. There is however, potential for significant impact from this pest for day neutral strawberries which continue fruiting when populations tend to increase. SWD appears to have the capability to survive winter conditions. However, populations at the start of the growing season tend to be quite low indicating high mortality over the winter. Adult flies are 2-3 mm in length, with red eyes and a tan-colored body with darker bands on the abdomen. Males have characteristic single spots at the leading edge of the tip of the wing and two dark spots on their front legs. Females lack wing spots and leg spots, but are distinguished by a saw-toothed ovipositor (visible under magnification). Larvae are white, nondescript and legless. Female SWD can lay eggs in ripening and marketable fruit.

Monitoring can be important for managing this pest. Talk to your local extension educator about a monitoring program. Traps and baits are now commercially available. Or homemade traps and baits, based on a fermenting mixture of yeast, sugar, water, and whole wheat flour with an apple cider vinegar drawing solution can be constructed (see <http://www.fruit.cornell.edu/spottedwing/> for more information). Fruit should also be inspected for evidence of larval feeding (see below).

Fruit destined for a processing market may be at risk of rejection due to presence of larvae. Home canning and processing may generate complaints from customers that notice SWD larvae. **Maintain a good cold chain between harvest and sale.** Display farm market fruit in a cooler—refrigeration slows or stops SWD development in fruit. Regular fruit sampling will help identify problems in the field. Fruit can be inspected for evidence of larval feeding. Small pinholes in berries may leak juice when the berry is gently squeezed. Immersing fruit in a salt solution (1 Tbsp. table salt/cup water (14.8 cc/236.6 ml)) may cause larvae to float to surface. At least 100 fruit per block per harvest should be observed for infestation.

For more information, consult the [Spotted Wing Drosophila website](#) on Cornell Fruit Resources.

Spotted Wing Drosophila Management Options	
Scouting/thresholds	Not specifically established but customer tolerance for infested fruit is likely to be very low.
Variety susceptibility	No known resistant varieties.
Cultural management	<p>Canopy and water management will make the environment less favorable. Use a adequate plant and rows pacing at planting to maintain an open canopy, increase sunlight and reduce humidity. Similarly, at renovation, narrow June-bearing strawberry matted rows to an 8 to 10" width. These practices will make plantings less attractive to SWD and will improve spray coverage. Repair leaking driplines and avoid overhead irrigation when possible. Allow the ground and mulch surfaces to dry before irrigating.</p> <p>Excellent sanitation will reduce SWD populations. Fruit should be harvested frequently and completely to prevent the buildup of ripe and over-ripe fruit. Unmarketable fruit should be removed from the field and either frozen, "baked" in clear plastic bags placed in the sun, or disposed of in bags off-site. This will kill larvae, remove them from your crop, and prevent them from emerging as adults.</p> <p>Cool berries immediately. Chilling berries immediately after harvest to 32-33F will slow or stop the development of larvae and eggs in the fruit. U-Pick customers should be encouraged to follow this strategy to improve fruit quality at home.</p> <p>If the planting includes day neutral varieties; consider using insect exclusion netting on these to protect fruit; if establishing a new planting, focus on June-bearing varieties to minimize the need for SWD management.</p>
Chemical treatment	A few insecticides have recently been granted 2ee label exemptions for control of SWD. SWD adults appear sensitive to several different chemistries, although their high reproductive rate, short generation time, and mobility may necessitate multiple applications for control.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.6.5 Pesticides Labeled for Management of Spotted Wing Drosophila					
Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Aza-Direct (azadirachtin)	1-2 pts/acre	0	4	2	
AzaGuard (azadirachtin)	10-16 fl oz/acre	0	4	2	Apply with OMRI approved spray oil.
AzaMax (azadirachtin)	1.33 fl oz/1000 sq ft	0	4	2	
AzaSol (azadirachtin)	6 oz/acre	-	4	2	Spray when larvae first appear.
Azatrol-EC (azadirachtin)	0.24-0.96 fl oz/1000 sq ft	0	4	2	
Azera (azadirachtin, pyrethrins)	1-3.5 pts/acre	-	12	1	
Ecozin Plus 1.2% ME (azadirachtin)	15-30 oz/acre	0	4	?	
Entrust (spinosad)	1.25-2 oz/acre	1	4	1	2(ee) recommendation. User must have a

Table 7.6.5 Pesticides Labeled for Management of Spotted Wing Drosophila					
Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
					copy of the recommendation in their possession at the time of application. http://pims.psur.cornell.edu/LabelResults.php?ProductId=154869&SearchPage=ProductName.php
Entrust SC (spinosad)	4-6 fl oz/acre	1	4	1	2(ee) recommendation. User must have a copy of the recommendation in their possession at the time of application. http://pims.psur.cornell.edu/LabelResults.php?ProductId=176736&SearchPage=ProductName.php
Envirepel 20 (garlic juice)	10-32 oz/acre	-	-	?	25(b) pesticide. Repellant.
Molt-X (azadirachtin)	10 oz/acre	0	4	2	
PyGanic EC 1.4 II (pyrethrins)	16-64 fl oz/acre	until dry	12	2	Short residual activity may require multiple applications. Caution: do not use when bees are active in the planting.
PyGanic EC 5.0 II (pyrethrins)	4.5-17 fl oz/acre	until dry	12	2	See note on PyGanic EC 1.4 II

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.

PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.7 MINOR AND SPORADIC INSECT AND MITE PESTS

Many insects found in strawberry plantings of New York, while having the capacity to cause economic damage, do not occur on a yearly basis at damaging levels and therefore are considered minor or sporadic pests. For these reasons it is important to be familiar with the life cycle of the pest to assist in developing a scouting program that will ensure a pest problem can be discovered and dealt with before it becomes an outbreak. And again, it is important to know when a potential pest is not causing significant economic damage so that unnecessary controls can be avoided.

7.7.1 BUD WEEVIL (CLIPPER) (*Anthonomus signatus*)

Adults puncture blossom buds while feeding in the spring, deposit eggs in the nearly mature buds, and then girdle the bud so that it hangs by a mere thread or falls to the ground. Injury is most likely along edges of fields or when strawberries are grown next to woodlots or other sites suitable for adult hibernation. Frequent scouting for bud cutting is important in areas where weevil pressure is expected to be high. In the past, a treatment threshold of 1 cut bud per linear foot has been recommended. Research conducted in the last few years, however, suggests that plants can sustain many times this pressure without a measurable reduction in yield if dipping occurs on tertiary flower buds. The new threshold is more than one primary or secondary flower bud or more than two tertiary flower buds per truss, or more than one injured truss per foot of row. Mulches and full-canopy beds may encourage newly emerged adults to remain in the planting so that damage increases in succeeding years. Using cropping systems shorter than 3 years, plowing under all old beds immediately after final harvest, and removing foliage and mulch to reduce the suitability of overwintering sites help lessen the chances of dipper injury.

IPM fact sheet Bud Weevil (Clipper) nysipm.cornell.edu/factsheets/berries/strawberry_dipper.pdf

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.7.1 Pesticides Labeled for Management of Bud Weevil

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Aza-Direct (azadirachtin)	1-2 pts/acre	0	4	?	Foliar spray or soil drench.
AzaGuard (azadirachtin)	10-16 fl oz/acre	0	4	?	Apply with OMRI approved spray oil. Foliar spray or soil drench.
AzaMax (azadirachtin)	1.33 fl oz/1000 ft ²	0	4	?	Foliar spray or soil drench.
AzaSol (azadirachtin)	6 oz/acre	-	4	?	Foliar spray or soil drench.
Azatrol-EC (azadirachtin)	0.29-0.96 fl oz/1000 ft ²	0	4	?	
Azera (azadirachtin, pyrethrins)	1-3.5 pts/acre	-	12	?	
BioLink (garlic juice)	0.5-2 qt/acre	-	-	?	25(b) pesticide.
BioLink Insect & Bird Repellent (garlic juice)	0.5-4 qt/acre	-	-	?	25(b) pesticide.
Ecozin Plus 1.2% ME (azadirachtin)	15-30 oz/acre	0	4	?	
Envirepel 20 (garlic juice)	10-32 oz/acre	-	-	?	25(b) pesticide. Repellent.
Garlic Barrier AG (garlic juice)	See comments	-	-	?	25(b) pesticide. See label for specific information.
Molt-X (azadirachtin)	10 oz/acre	0	4	?	
PyGanic EC 1.4 II (pyrethrins)	16-64 fl oz/acre	until dry	12	?	Spraying should begin when the insects first appear. Repeat as required. Caution: do not use when bees are active in the planting.
PyGanic EC 5.0 II (pyrethrins)	4.5-17 fl oz/acre foliar spray	until dry	12	?	Short residual activity may require multiple applications. Caution: do not use when bees are active in the planting.
Safer Brand #567 II (potassium laurate, pyrethrins)	6.4 oz/ gal water	until dry	12	?	

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.

PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.7.2 SPITTLEBUG (*Philaenus spumaris*)

White frothy masses on the stems and leaves around the time of bloom harbor the nymphs, which pierce the stems and suck plant juices. Their feeding, if extensive, can stunt the plants and reduce berry size. Leaves appear crinkled and darker green than undamaged leaves. The spittle masses are a great nuisance to pickers. Threshold is one mass per square ft of row. Good weed control may help to reduce numbers. Populations are usually largest in weedy fields. Only one generation is produced per year. The leaves recover after the insects are gone.

IPM fact sheet Spittlebug nysipm.cornell.edu/factsheets/berries/meadow_spittlebug.pdf

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.7.2 Pesticides Labeled for Management of Spittlebug

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Aza-Direct (azadirachtin)	1-2 pts/acre	0	4	?	
AzaGuard (azadirachtin)	10-16 fl.oz./acre	0	4	?	Apply with OMRI approved spray oil.
AzaMax (azadirachtin)	1.33 fl oz/1000 ft ²	0	4	?	
AzaSol (azadirachtin)	6 oz/ 50 gal water	-	4	?	For nymph treatment.
Azatrol-EC (azadirachtin)	0.24-0.96 fl oz/1000 ft ²	0	4	?	
Azera (azadirachtin, pyrethrins)	1-3.5 pts/acre	-	12	?	
Ecozin Plus 1.2% ME (azadirachtin)	15-30 oz/acre	0	4	?	
Envirepel 20 (garlic juice)	10-32 oz/acre	-	-	?	25(b) pesticide. Repellent.
Garlic Barrier AG (garlic juice)	See comments.	-	-	?	25(b) pesticide. See label for specific information.
Molt-X (azadirachtin)	10 oz/acre	0	4	?	
Neemix 4.5 (azadirachtin)	7-16 fl oz/acre	0	4	?	
PyGanic EC 1.4 II (pyrethrins)	16-64 fl oz/acre	until Dry	12	1	

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.

PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.7.3 STRAWBERRY ROOTWORM (*Paria fragariae-complex*)

Larvae feed on roots in late spring to early summer. Adults feed on leaves in May and again in late July, at night.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.7.3 Pesticides Labeled for Management of Strawberry Rootworm

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Aza-Direct (azadirachtin)	1-2 pts/acre	0	4	3	Foliar spray or soil drench.
AzaGuard (azadirachtin)	8-16 fl oz/acre	0	4	3	Apply with OMRI approved spray oil. Foliar spray or soil drench.
AzaMax (azadirachtin)	1.33 fl oz/1000 ft ²	0	4	3	Foliar spray or soil drench.
AzaSol (azadirachtin)	6 oz/ 50 gal water	-	4	3	Foliar spray or soil drench.
Azatrol-EC (azadirachtin)	0.29-0.96 fl oz/1000 ft ²	0	4	3	
Azera (azadirachtin, pyrethrins)	1-3.5 pts/acre	-	12	3	Foliar spray or soil drench.
Envirepel 20 (garlic juice)	10-32 oz/acre	-	-	?	25(b) pesticide. Repellent.

Table 7.7.3 Pesticides Labeled for Management of Strawberry Root tworm

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Garlic Barrier AG (garlic juice)	See comments	-	-	?	25(b) pesticide. See label for specific information.
Molt-X (azadirachtin)	8 oz/acre	0	4	3	Foliar spray or soil drench.
PyGanic EC 1.4 II (pyrethrins)	16-64 fl oz/acre	until dry	12	?	Spraying should begin when the insects first appear. Repeat as required. Caution: do not use when bees are active in the planting.
PyGanic EC 5.0 II (pyrethrins)	4.5-17 fl oz/acre	until dry	12	?	See note on PyGanic EC 1.4 II

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.

PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.7.4 GREENHOUSE WHITEFLY (*Trialeurodes vaporariorum*)

Whiteflies are small, white insects that resemble flies but are actually more closely related to aphids. Whiteflies feed on young plants, causing stunting.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.7.4 Pesticides Labeled for Management of Greenhouse Whitefly

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Aza-Direct (azadirachtin)	1-2 pts/acre	0	4	?	
AzaGuard (azadirachtin)	8-21 fl.oz./acre	0	4	?	Apply with OMRI approved spray oil.
AzaMax (azadirachtin)	1.33 fl oz/1000 ft ²	0	4	?	
AzaSol (azadirachtin)	6 oz/ 50 gal water	-	4	?	
Azatrol-EC (azadirachtin)	0.24-0.96 fl oz/1000 ft ²	0	4	?	
Azera (azadirachtin, pyrethrins)	1-3.5 pts/acre	-	12	?	
BioLink (garlic juice)	0.5-2 qt/acre	-	-	?	25(b) pesticide.
BioLink Insect & Bird Repellent (garlic juice)	0.5-4 qt/acre	-	-	?	25(b) pesticide.
BioRepel (garlic oil)	1 part BioRepel to 100 parts water	-	-	?	25(b) pesticide.
Cedar Gard (cedar oil)	1 qt/acre	-	-	?	25(b) pesticide.
Ecotec (rosemary oil, peppermint oil)	1-4 pt/acre	-	-	3	25(b) pesticide.
Ecozin Plus 1.2% ME (azadirachtin)	15-30 oz/acre	0	4	?	
Envirepel 20 (garlic juice)	10-32 oz/acre	-	-	?	25(b) pesticide. Repellent.
Garlic Barrier AG (garlic juice)	See comments.	-	-	?	25(b) pesticide. See label for specific information.

Table 7.7.4 Pesticides Labeled for Management of Greenhouse Whitefly

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Grandevo (Chromobacterium subtsugae str. PR4A4-1)	2-3 lb/acre	0	4	?	
Molt-X (azadirachtin)	8 oz/acre	0	4	?	
M-Pede (insecticidal soap)	1-2% vol/vol	0	12	?	Works by contact. Good coverage is important. Use in combination with another pesticide for enhanced and residual effect.
Neemix 4.5 (azadirachtin)	4-16 fl oz/acre	0	4	?	Rate and frequency of application vary based on pest pressure. See label for guidance.
Nuke Em (citric acid)	1 fl oz/31 fl oz water. Normal strength.	-	-	?	25(b) pesticide. Use the normal strength mix first. See label for stronger dilutions if needed.
Oleotrol-I Bio-Insecticide Concentrate (soybean oil)	43-45 fl oz/100 gal water	-	-	?	25(b) pesticide.
Organocide (sesame oil)	1-2 gal/100 gal water	-	-	?	25(b) pesticide.
PFR-97 20% WDG (Isaria fumosorosea Apopka str. 97)	1-2 lb/acre	-	4	?	
PureSpray Green (white mineral oil)	0.75-1.5 gal/100 gal water	up to day	4	?	Spray at no less than 400 PSI using ceramic nozzles.
PyGanic EC 1.4 II (pyrethrins)	16-64 fl.oz./acre	until dry	12	?	Spraying should begin when the insects first appear. Repeat as required. Caution: do not use when bees are active in the planting.
PyGanic EC 5.0 II (pyrethrins)	4.5-17 fl.oz./acre	until dry	12	?	See comment for PyGanic EC 1.4 II
Sil-Matrix (potassium silicate)	0.5-1% solution	0	4	?	
SuffOil-X (aliphatic petroleum solvent)	1-2 gal/100 gal water	up to day	4	?	Can cause phytotoxicity if applied too close to a sulfur application.
Trilogy (neem oil)	1-2% solution	up to day	4	?	Maximum labeled use of 2 gal/acre/application. Apply in sufficient water to achieve complete coverage. Provides suppression only.
TriTek (mineral oil)	1-2 gal/100 gal water	up to day	4	?	Apply as needed.

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.

PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.7.5 CYCLAMEN MITE (*Stenotarsonemus pallidus*)

This tiny (one one-hundredth-inch) mite is pinkish orange and shiny when mature. Its translucent eggs are often so abundant that they appear as a white mass along the mid-veins of folded, newly emerging leaves. The mites feed on the young leaves in plant crowns; when the leaves emerge, they are stunted, crinkled, and malformed. Blossom feeding later

results in misshapen fruit. The mites are most troublesome in strawberry beds that are kept for long periods, although in some cases young plantings will have them. They increase in number during bloom and peak during fruit development. Avoid infested planting stock. ‘Cabot’ is particularly susceptible. Insectary-reared predatory mites may provide some control of cyclamen mites. *Neoseiulus fallacis* and *N. cucumeris* have been found to provide some control by researchers.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide’s effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIIMS website](#)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.7.5 Pesticides Labeled for Management of Cyclamen Mite

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Aza-Direct (azadirachtin)	1-2 pts/acre	0	4	1	
AzaGuard (azadirachtin)	10-16 fl oz/acre	0	4	1	Apply with OMRI approved spray oil.
AzaMax (azadirachtin)	1.33 fl oz/1000 ft ²	0	4	1	
Azatrol-EC (azadirachtin)	0.24-0.96 fl oz/1000 ft ²	0	4	1	
BioLink (garlic juice)	0.5-2 qt/acre	-	-	?	25(b) pesticide.
BioLink Insect & Bird Repellent (garlic juice)	0.5-4 qt/acre	-	-	?	25(b) pesticide.
Ecotec (rosemary oil, peppermint oil)	1-4 pt/acre	-	-	?	25(b) pesticide.
Envirepel 20 (garlic juice)	10-32 oz/acre	-	-	?	25(b) pesticide. Repellent.
Garlic Barrier AG (garlic juice)	See comments	-	-	?	25(b) pesticide. See label for specific information.
GC-Mite (garlic oil, clove oil, cottonseed oil)	1 gal/100 gal water	-	-	1	25(b) pesticide. Conduct compatibility test prior to application.
Grandevo (<i>Chromobacterium subtsugae</i> str. PRAA4-1)	2-3 lb/acre	0	4	?	
JMS Stylet-Oil (paraffinic oil)	3 qt/100 gal water	0	4	2	Apply for optimum coverage of leaf surfaces. Use high pressure, small droplet size, and adequate gallonage to ensure good coverage. Can cause phytotoxicity if applied too close to a sulfur application.
Nuke Em (citric acid)	1 fl oz/ 31 fl oz water. Normal strength	-	-	?	25(b) pesticide. Use the normal strength mix first. See label for stronger dilutions if needed.
Oleotrol-I Bio-Insecticide Concentrate (soybean oil)	43-45 fl oz/100 gal water	-	-	?	25(b) pesticide.
PureSpray Green (white mineral oil)	0.75-1.5 gal/100 gal water	up to day	4	1	Spray at no less than 400 PSI using ceramic nozzles.
PyGanic EC 1.4 II (pyrethrins)	16-64 fl oz/acre	until dry	12	?	Short residual activity may require multiple applications. Caution: do not use when bees

Table 7.7.5 Pesticides Labeled for Management of Cyclamen Mite

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
					are active in the planting.
PyGanic EC 5.0 II (pyrethrins)	4.5-17 fl oz/acre	until dry	12	?	See comment on PyGanic EC 1.4 II
Sil-Matrix (potassium silicate)	0.5-1% solution	0	4	?	
SuffOil-X (aliphatic petroleum solvent)	1-2 gal/100 gal water	up to day	4	1	Can cause phytotoxicity if applied too close to a sulfur application.
Trilogy (neem oil)	1-2% solution	up to day	4	?	Maximum labeled use of 2 gal/acre/application
TriTek (mineral oil)	1-2 gal/100 gal water	up to day	4	1	Apply as needed.

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.

PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.7.6 LEAFROLLER (*various species*)

Several species of moth larvae roll or fold strawberry leaves with silk. Leaf injury can be seen throughout the season, but an extremely large population is required before noticeable crop damage occurs.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PPIMS website](#)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.7.6 Pesticides Labeled for Management of Leafrollers

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Aza-Direct (azadirachtin)	1-2 pts/acre	0	4	?	
AzaGuard (azadirachtin)	8-16 fl oz/acre	0	4	?	Apply with OMRI approved spray oil.
AzaMax (azadirachtin)	1.33 fl oz/1000 ft ²	0	4	?	
AzaSol (azadirachtin)	6 oz/ 50 gal water	-	4	?	
Azatrol-EC (azadirachtin)	0.24-0.96 fl oz/1000 ft ²	0	4	?	
Azera (azadirachtin, pyrethrins)	1-3.5 pts/acre	-	12	?	See label for specific leafroller species product can be used against.
Biobit HP (<i>Bacillus thuringiensis</i> subsp. <i>Kurstaki</i>)	0.5-1 lb/acre	0	4	1	See comment for Azera.
BioLink (garlic juice)	0.5-2 qt/acre	-	-	?	25(b) pesticide.
BioLink Insect & Bird Repellent (garlic juice)	0.5-4 qt/acre	-	-	?	25(b) pesticide.
Cedar Gard (cedar oil)	1 qt/acre	-	-	?	25(b) pesticide.
Deliver (<i>Bacillus thuringiensis</i> subsp. <i>Kurstaki</i>)	0.25-1.5 lb/acre	0	4	1	See comment for Azera.

Table 7.7.6 Pesticides Labeled for Management of Leafrollers

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Dipel DF (<i>Bacillus thuringiensis</i> subsp. <i>Kurstaki</i>)	0.5-1 lb/acre	0	4	1	See comment for Azera.
Ecotec (rosemary oil, peppermint oil)	1-4 pts/acre	-	-	?	25(b) pesticide.
Ecozin Plus 1.2% ME (azadirachtin)	15-30 oz/acre	0	4	?	
Entrust (spinosad)	1.25-2 oz/acre	1	4	1	Treat when pests appear, targeting eggs at hatch or small larvae.
Entrust SC (spinosad)	4-6 fl oz/acre	1	4	1	See comment for Entrust.
Envirepel 20 (garlic juice)	10-32 oz/acre	-	-	?	25(b) pesticide. Repellent.
Garlic Barrier AG (garlic juice)	See comments	-	-	?	25(b) pesticide. See label for specific information.
Grandevo (<i>Chromobacterium subtsugae</i> str. PRAA4-1)	1-3 lb/acre	0	4	?	
Javelin WG (<i>Bacillus thuringiensis</i> subsp. <i>Kurstaki</i>)	0.25-1.5 lb/acre	0	4	1	See comment for Azera.
Molt-X (azadirachtin)	8 oz/acre	0	4	?	
Neemix 4.5 (azadirachtin)	7-16 fl oz/acre	0	4	?	See comment for Azera.
Organocide (sesame oil)	1-2 gal/100 gal water	-	-	?	25(b) pesticide.
PyGanic EC 1.4 II (pyrethrins)	16-64 fl oz/acre	until dry	12	?	Spraying should begin when the insects first appear. Repeat as required. Caution: do not use when bees are active in the planting.
PyGanic EC 5.0 II (pyrethrins)	4.5-17 fl.oz./acre	until dry	12	?	See comment for PyGanic EC 1.4 II
XenTari (<i>Bacillus thuringiensis</i> , var. <i>aizawai</i>)	0.5-1.5 lb/acre	0	4	1	See comment for Azera

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.

PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.7.7 APHIDS (*various species*)

These soft-bodied insects usually occur on new shoots and buds in the crown of the plant and along the veins on the undersides of the leaves. When present in large numbers, they weaken the plant. They have also been associated with some viruses. Their honeydew promotes the growth of a black sooty mold, which makes the fruit and leaves sticky, hindering harvest and reducing marketability. More important, aphids are vectors for several serious virus diseases. Aphid populations often are controlled by natural enemies and do not require insecticide control.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.7.7 Pesticides Labeled for Management of Aphids

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Aza-Direct (azadirachtin)	1-2 pts/acre	0	4	1	
AzaGuard (azadirachtin)	10-16 fl oz/acre	0	4	1	Apply with OMRI approved spray oil.
AzaMax (azadirachtin)	1.33 fl oz/1000 ft ²	0	4	1	
AzaSol (azadirachtin)	6 oz/ 50 gal water	-	4	1	
Azatrol-EC (azadirachtin)	0.24-0.96 fl oz/1000 ft ²	0	4	1	
Azera (azadirachtin, pyrethrins)	1-3.5 pts/acre	-	12	1	
BioLink (garlic juice)	0.5-2 qt/acre	-	-	?	25(b) pesticide.
BioLink Insect & Bird Repellent (garlic juice)	0.5-4 qt/acre	-	-	?	25(b) pesticide.
BioRepel (garlic oil)	1 part BioRepel to 100 parts water	-	-	?	25(b) pesticide.
DES-X (insecticidal soap)	2% solution sprayed at 75-200 gallons/acre	1/2	12	?	
Ecotec (rosemary oil, peppermint oil)	1-4 pt/acre	-	-	?	25(b) pesticide.
Ecozin Plus 1.2% ME (azadirachtin)	15-30 oz/acre	0	4	1	
Envirepel 20 (garlic juice)	10-32 oz/acre	-	-	?	25(b) pesticide. Repellent.
Garlic Barrier AG (garlic juice)	See comments	-	-	?	25(b) pesticide. See label for specific information.
GC-Mite (garlic oil, clove oil, cottonseed oil)	1 gal/100 gal water	-	-	1	25(b) pesticide. Conduct compatibility test prior to application.
Grandevo (<i>Chromobacterium subtsugae</i> str. PRAA4-1)	2-3 lb/acre	0	4	?	
GrasRoots (cinnamon oil)	1 part GrasRoots: 9 parts water	-	-	?	25(b) pesticide.
Molt-X (azadirachtin)	10 oz/acre	0	4	1	
M-Pede (insecticidal soap)	1-2% vol/vol	0	12	1	Works by contact. Good coverage is important.
Neemix 4.5 (azadirachtin)	5-7 fl oz/acre	0	4	1	
Nuke Em (citric acid)	1 fl oz/ 31 fl oz water. Normal strength.	-	-	?	25(b) pesticide. Use the normal strength mix first. See label for stronger dilutions if needed.

Table 7.7.7 Pesticides Labeled for Management of Aphids

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Oleotrol-I Bio-Insecticide Concentrate (soybean oil)	43-45 fl oz/100 gal water	-	-	?	25(b) pesticide.
Organocide (sesame oil)	1-2 gal/100 gal water	-	-	1	25(b) pesticide.
PFR-97 20% WDG (Isaria fumosorosea Apopka str. 97)	1-2 lb/acre	-	4	?	
PureSpray Green (white mineral oil)	0.75-1.5 gal/100 gal water	up to day	4	1	Spray at no less than 400 PSI using ceramic nozzles.
PyGanic EC 1.4 II (pyrethrins)	16-64 fl oz /acre	until dry	12	2	Spraying should begin when the insects first appear. Repeat as required. Caution: do not use when bees are active in the planting.
PyGanic EC 5.0 II (pyrethrins)	4.5-17 fl oz./acre	until dry	12	2	See comment for PyGanic EC 1.4 II
Safer Brand #567 II (potassium laurate, pyrethrins)	6.4 oz/ gal water	until dry	12	?	
Sil-Matrix (potassium silicate)	0.5-1% solution.	0	4	?	
SuffOil-X (aliphatic petroleum solvent)	1-2 gal/100 gal water	up to day	4	1	Can cause phytotoxicity if applied too close to a sulfur application.
Trilogy (neem oil)	1-2% solution	up to day	4	?	Maximum labeled use of 2 gal/acre/application. Apply in sufficient water to achieve complete coverage.
TriTek (mineral oil)	1-2 gal/100 gal water	up to day	4	1	Apply as needed.

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.

PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.7.8 POTATO LEAFHOPPER (*Empoasca fabae*)

Adults migrate into New York State in early to mid-June, carried by summer weather fronts. The adults and nymphs feed along the veins on the undersides of leaves by sucking plant juices, and in the process, inject a toxic substance with their saliva. Affected plants have shortened petioles and small distorted leaves that bend down at right angles. Leaf yellowing is also seen, starting at the margins and progressing toward the mid-vein. Avoid proximity to alfalfa plantings, which provide a major source of potato leafhopper population build-up.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PLIMS website](#)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.7.8 Pesticides Labeled for Management of Potato Leafhopper

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Aza-Direct (azadirachtin)	1-2 pts/acre	0	4	1	
AzaGuard (azadirachtin)	10-16 fl oz/acre	0	4	1	Apply with OMRI approved spray oil.
AzaMax (azadirachtin)	1.33 fl oz/1000 sq ft	0	4	1	
Azatrol-EC (azadirachtin)	0.24-0.96 fl oz/1000 sq ft	0	4	1	
Azera (azadirachtin, pyrethrins)	1-3.5 pts/acre	-	12	1	
BioLink (garlic juice)	0.5-2 qt/acre	-	-	?	25(b) pesticide.
BioLink Insect & Bird Repellent (garlic juice)	0.5-4 qt/acre	-	-	?	25(b) pesticide.
BioRepel (garlic oil)	1 part BioRepel to 100 parts water	-	-	?	25(b) pesticide.
Cedar Gard (cedar oil)	1 qt/acre	-	-	?	25(b) pesticide.
DES-X (insecticidal soap)	2% solution sprayed at 75-200 gallons/acre	1/2	12	?	
Ecotec (rosemary oil, peppermint oil)	1-4 pt/acre	-	-	?	25(b) pesticide.
Ecozin Plus 1.2% ME (azadirachtin)	15-30 oz/acre	0	4	1	
Envirepel 20 (garlic juice)	10-32 oz/acre	-	-	?	25(b) pesticide. Repellent.
Garlic Barrier AG (garlic juice)	See comments	-	-	?	25(b) pesticide. See label for specific information.
Molt-X (azadirachtin)	10 oz/acre	0	4	1	
M-Pede (insecticidal soap)	1-2% vol/vol	0	12	3	Works by contact. Good coverage is important.
Neemix 4.5 (azadirachtin)	7-16 fl oz/acre	0	4	1	
PyGanic EC 1.4 II (pyrethrins)	16-64 fl oz/acre	until dry	12	1	Spraying should begin when the insects first appear. Repeat as required. Caution: do not use when bees are active in the planting.
PyGanic EC 5.0 II (pyrethrins)	4.5-17 fl oz/acre	until dry	12	1	See comment on PyGanic EC 1.4 II
Safer Brand #567 II (potassium laurate, pyrethrins)	6.4 oz/ gal water	until dry	12	?	

Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.

PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.7.9 JAPANESE BEETLE (*Popillia japonica*)

Beetles emerge in early July and feed on leaves. Although there are Japanese beetle traps, research has shown that the traps may attract more beetles into a planting than they eliminate in the traps.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PLMS website](https://www.dec.state.ny.us/da/pubs/plms/)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.7.9 Pesticides Labeled for Management of Japanese Beetle

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
Aza-Direct (azadirachtin)	1-2 pts/acre	0	4	3	
AzaGuard (azadirachtin)	8-16 fl oz/acre	0	4	3	Apply with OMRI approved spray oil.
AzaMax (azadirachtin)	1.33 fl oz/1000 ft ²	0	4	3	
AzaSol (azadirachtin)	6 oz/ 50 gal water	-	4	3	Spray when larvae first appear.
Azatrol-EC (azadirachtin)	0.29-0.96 fl oz/1000 ft ²	0	4	3	
Azera (azadirachtin, pyrethrins)	1-3.5 pts/acre	-	12	3	
BioLink (garlic juice)	0.5-2 qt/acre	-	-	?	25(b) pesticide.
BioLink Insect & Bird Repellent (garlic juice)	0.5-4 qt/acre	-	-	?	25(b) pesticide.
Cedar Gard (cedar oil)	1 qt/acre	-	-	?	25(b) pesticide.
Ecozin Plus 1.2% ME (azadirachtin)	15-30 oz/acre	0	4	3	
Envirepel 20 (garlic juice)	10-32 oz/acre	-	-	?	25(b) pesticide. Repellent.
Garlic Barrier AG (garlic juice)	See comments	-	-	?	25(b) pesticide. See label for specific information.
Molt-X (azadirachtin)	8 oz/acre	0	4	3	
PyGanic EC 1.4 II (pyrethrins)	16-64 fl oz/acre	until dry	12	3	Spraying should begin when the insects first appear. Repeat as required. Caution: do not use when bees are active in the planting.
PyGanic EC 5.0 II (pyrethrins)	4.5-17 fl oz/acre	until dry	12	3	See comment for PyGanic EC 1.4 II
Safer Brand #567 II (potassium laurate, pyrethrins)	6.4 oz/ gal water	until dry	12	?	

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.

PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.8 Slug Management (*various species*)

These soft-bodied mollusks resemble snails without a shell. Slugs feed on ripening fruit, leaving holes in the berries. They are most active at night and during cool, wet weather. Populations are greatest when the weather is damp and the planting is mulched. Translucent silver to whitish slime trails are visible on damaged plant parts.

IPM fact sheet Banded Slug nysipm.cornell.edu/factsheets/fieldcrops/b_slug.pdf

IPM fact sheet Gray Garden Slug nysipm.cornell.edu/factsheets/fieldcrops/gg_slug.pdf

IPM fact sheet Marsh Slug nysipm.cornell.edu/factsheets/fieldcrops/m_slug.pdf

IPM fact sheet Spotted Garden Slug nysipm.cornell.edu/factsheets/fieldcrops/sg_slug.pdf

Slug Management Options	
Scouting/thresholds	None established.
Variety susceptibility	No known resistant varieties.
Cultural management	<p>Eliminating mulch will reduce slug populations, but will cause other problems, so this is not recommended.</p> <p>Good sanitation and weed control helps to reduce slug populations.</p> <p>In areas where slugs are a problem, avoid perennial clovers as cover crops and rotate out of alfalfa or other perennial legumes 1 year prior to planting establishment.</p> <p>Overhead irrigation creates conditions especially favorable to slugs. If overhead irrigation must be used, irrigate during morning hours to allow foliage to dry before evening.</p>
Chemical treatment	See below.

At the time this guide was produced, the following materials were available in New York State for managing this pest and were allowable for organic production. Listing a pest on a pesticide label does not assure the pesticide's effectiveness. The registration status of pesticides can and does change. Pesticides must be currently registered with the New York State Department of Environmental Conservation (DEC) to be used legally in NY. However, pesticides meeting the federal requirements for minimum-risk (25(b)) pesticides do not require registration. Current NY pesticide registrations can be checked on the Pesticide Product, Ingredient, and Manufacturer System ([PIMS website](#)). ALWAYS CHECK WITH YOUR CERTIFIER before using a new product.

Table 7.8.1 Pesticides Labeled for Management of Slugs

Trade Name (active ingredient)	Product Rate	PHI (Days)	REI (Hours)	Efficacy ¹	Comments
BioLink (garlic juice)	0.5-2 qt/acre	-	-	?	25(b) pesticide.
BioLink Insect & Bird Repellent (garlic juice)	0.5-4 qt/acre	-	-	?	25(b) pesticide.
Bug-N-Sluggo (spinosad, iron phosphate)	20-44 lb/acre soil treatment	1	4	?	
Garlic Barrier AG (garlic juice)	See comments	-	-	?	25(b) pesticide. See label for specific information.
Sluggo-AG (iron phosphate)	20-44 lb/acre soil treatment	0	0	?	Spread bait around perimeter to intercept slugs migrating toward berries.
Sluggo Slug and Snail Bait (iron phosphate)	20-44 lb/acre soil treatment	0	0	?	

¹ Efficacy: 1-effective in some research studies, 2- inconsistent efficacy results, 3-not effective, ?-not reviewed or no research available.

PHI-Pre-harvest interval. REI-Restricted Entry Interval. - = pre-harvest interval isn't specified on label.

7.9 Wildlife Management

Various rodents can damage a strawberry planting, especially as they feed under mulch in the winter. Closely mowing the area around the planting in early November will reduce the habitat for voles and mice. The habitats (woodlots) of predators that feed on rodents (hawks, owls, foxes) should be protected around the area. A number of toxic baits are labeled for use in agricultural areas. To be most effective, baits should be placed in feeding stations that exclude large animals and are replenished throughout the winter.

Deer browsing can devastate berry plantings. Multiple strategies are required to discourage deer from feeding on berry plantings. Refer to [Reducing Deer Damage to Home Gardens and Landscape Plantings](#), by P. Curtis and M. Richmond for recommended methods. Fencing is the best way to keep deer and other mammals out of berry plantings. Some deer repellents are registered for use on fruit crops during the non-bearing season.

When using dogs and invisible fence to manage vertebrate pests in a planting, there is food safety risk associated with the dog excrement. If the dog consistently excretes in an area away from the field, or keeps other vertebrate animals from using the field, the food safety risk is somewhat reduced. Using dogs primarily in the winter and early spring when deer browsing is greatest (and avoiding use during harvest) will also minimize food safety risk.

Table 7.9.1. Vertebrate Damage Mitigation Practices

Animal Pest	Management Practices¹
Mice and voles	Removal of dropped fruit; habitat manipulations including elimination of unmowable areas surrounding plantings; monitor to determine the need for management. Mow closely in late fall around the planting and apply winter mulch only after mowing. Population control through trapping by landowner.
Raccoons	Avoid sites with woods along the edge(s) because these will support raccoon populations. Electrified exclusion fencing. Population reduction through shooting by licensed hunters or landowners in appropriate seasons; through trapping by landowner, by licensed trapper, or by licensed nuisance wildlife control operator.
Red and gray foxes	Tend to chew on irrigation lines. Manipulation including elimination of protective cover around plantings. Population reduction through shooting by licensed hunters or landowners in appropriate seasons; through trapping by landowner, by licensed trapper, or by licensed nuisance wildlife control operator.
White-tailed deer	Exclusion fencing (8 ft. [250 cm] high-tensile woven wire or 5 to 6 ft. [150 to 200 cm] electric exclusion fencing; peanut-butter baited electric fences; invisible fencing with dogs); habitat manipulation including elimination of protective cover around plantings. Population reduction through shooting by licensed hunters, landowners or their agents with DMAP or deer damage permits. Unlike with other vertebrate pests, landowners cannot kill nuisance deer without a permit.
Woodchucks	Exclusion fencing (electrified exclusion fencing); habitat manipulation including removal of brush piles. Population reduction through shooting by licensed hunters or landowners; through trapping by landowner or by licensed nuisance wildlife control operator.

¹ Conduct shooting and trapping only as defined by New York State Department of Environmental Conservation regulations. Shooting for nuisance wildlife control is allowed only when neighboring occupied buildings are >500 ft. distant; shooting when neighboring buildings are less than 500 ft. distant requires neighbor permission. Shooting also may require a permit, depending on animal and season. Also check local ordinances, as shooting and trapping are prohibited in some areas. Note: It is illegal to trap a nuisance animal and release it onto public lands or someone else's property. It must be released on the landowner's property or killed.

7.10 Considerations During Harvest and Renovation

During harvest operations some pests can become a nuisance, e.g. wasps and yellow jackets, particularly in U-pick operations. Wasp and yellow jacket nests can be destroyed during the growing season as they are found in the planting and surrounding areas. Some species are ground-nesting and such nests can be destroyed by drenching with hot water. Traps baited with sugary liquids, such as Hi-C, provide a means of reducing the population of wasps and yellow jackets, but the effectiveness of this tactic is unknown. For more information see "[Bee and Wasp Management, A Common Sense Approach](#)" (2011) by Jody Gangloff-Kaufman.

During harvest much can be done to reduce disease and insect pressure by eliminating infested and infected fruit from the planting. Separate damaged fruit from healthy fruit as it is being picked. Designate pickers to cull such fruit from the field at harvest time. Then bury or burn the diseased and infested fruit. This is helpful to combat gray mold, leather rot and anthracnose (through the removal of overripe and infected fruit), spotted wing drosophila, strawberry sap beetle (through the removal of overripe or infested fruit), and slugs (through the removal of overripe and dropped fruit).

After harvest, a post-harvest grading table will provide an excellent opportunity to grade out damaged, diseased and infected fruit which will lower quality and market value. All culled fruit should be destroyed by burning or burying. Cleanliness or sanitation in the planting is very important, removing dropped berries during harvest will reduce risk from gray mold, leather rot and anthracnose, spotted wing drosophila, strawberry sap beetle and slugs, as described above. At

this time also make note of trouble spots in the field, or the presence of unthrifty plants, foliar diseases, leaf damage, etc and plan steps to maintain a healthy planting.

At renovation do a thorough job of mowing the leaves off June-bearing strawberry plants, chopping mulch, and turning under infected and infested plant parts. Application of a thick mat of straw in early winter will provide protection from cold weather for the winter and assist in protecting plants from rain splashed inoculum from buried plant debris.

Keep in mind your production goals and recognize that it should be possible to obtain good yields in organic strawberry production. Therefore, maintain good records of the planting condition, pest pressure, the amount of fruit harvested, and your markets.

8. FOOD SAFETY

Attention to microbial food safety is important for crops that are eaten raw. Continuing produce-associated foodborne illness outbreaks have resulted in many buyers requiring the implementation of food safety practices on the farm and the development of the first ever produce safety regulations as part of the Food Safety Modernization Act (FSMA). Pathogens can contaminate food during all phases of production, harvesting and packing. Wild and domesticated animals, manure, irrigation water, inadequate worker hygiene, unclean picking containers, unsanitized post-harvest water, and unclean packaging materials are all potential vectors of microbiological contaminants. Growers should conduct a risk assessment to identify microbial hazards and then implement appropriate practices to reduce risks. There are many resources available to help including those at the [National GAPs Program](#) or the [Produce Safety Alliance](#). Regardless of farm size, commodities or cultural practices, Good Agricultural Practices can be used to identify and possibly reduce microbial risks.

Implementing just a few simple practices can reduce risks significantly. One of these is to wash hands using potable water and sanitizer prior to any contact with the crop, particularly after using the restroom or eating. Do not allow workers who are ill to handle produce. If they are able to work, assign jobs that do not involve contact with produce or customers. Prevent animals or animal manure from contacting produce, by discouraging animals (including pets) from entering production fields and by not using irrigation water that may have been contaminated with manure. Manure should only be applied before planting so it can be incorporated into the soil. For fall-fruited berries, composted manure can be applied to the soil in spring if it has been composted prior to application. Ensure that picking containers are clean and free from mouse droppings. Do not allow fruit to become wet after harvest. Following these steps can dramatically reduce risks of pathogen contamination. Conduct a full assessment of your farm to identify other high risk practices.

The Food Safety Modernization Act (FSMA) will apply to farms that grow, harvest, pack or hold most fruits and vegetables when those fruits and vegetables are in an unprocessed state, and will govern practices affecting: water, worker hygiene, manure and other soil additions, animals in the growing area, and equipment, tools and buildings. When the FSMA is finalized, the Food and Drug Administration (FDA) will be mandated to enforce preventive control measures, and to conduct inspections across the food supply system. Updates and information on this proposed rule are available at the United States Food and Drug Administration's [Food Safety Modernization Act](#) website.

Table 8.1 Rates for Sanitizers Labeled for Postharvest Strawberries and/or Postharvest Facilities			
Active ingredient Product name	Uses		
	Food contact surfaces¹	Hard surface, non-food contact¹	Fruit surface (spray or drench)
<i>chlorine dioxide</i>			
CDG Solution 3000	50 ppm solution	500 ppm dilution	5 ppm solution
Oxine ²	100 ppm solution	500 ppm solution	In tanks, use a 5 ppm solution; for process waters use a chemical feed pump or other injector system at 3 ¼ fl oz per 10 gal water. ³
Pro Oxine ²	50-200 ppm solution	500 ppm solution	-
<i>hydrogen peroxide/peroxyacetic acid</i>			
Enviroguard Sanitizer	-	2.5-20 fl oz/5 gal water	1 fl oz/20 gal water
Oxonia Active	1-1.4 oz/4 gal water	1 oz/8 gal water.	-
Peraclean 5	1-1.5 fl oz/5 gal water	-	-
Peraclean 15	0.33 fl oz/5 gal water	-	-
Perasan 'A'	1-6.1 oz/6 gal water	-	4z/20 gal water
Per-Ox	1-2.25 fl oz/5 gal water	1-10 fl oz/15 gal water	1 fl oz/5 gal water
SaniDate 5.0	1.6 fl oz/ 5 gal water	1.6 fl oz/ 5 gal water	59.1 to 209.5 fl oz/ 1,000 gallons water
SaniDate 12.0	-	-	25.6 to 89.6 fl oz / 1,000 gallons water
Shield-Brite PAA 5.0	1.6fl oz/5 gal water	1.6fl oz/5 gal water	59.1 to 209.5 fl. oz./1,000 gal water
Shield-Brite PAA 12.0	-	-	25.6 to 89.6 fl.oz/1,000 gal water
StorOx 2.0	0.5 fl oz/1 gal water	0.5 fl oz/1 gal water	1:220-1:1,000 dilution (processing/packing line treatment); 25-1.28 fl oz/100 gallons water (processing water); 0.35-0.58 fl oz/gallon (post-harvest spray)
Tsunami 100	-	-	2.5-6.7 fl oz/100 gal water
Victory	-	-	1 fl oz/16.4 gal water
VigorOx 15 F & V	0.31-0.45 fl oz/5 gal water-	1.1-9.5 fl oz/5 gal water -	0.54 fl oz/ 16 gal water
VigorOx LS-15	0.31-0.45 fl oz/5 gal water	1.1-9.5 fl oz/5 gal water	-
<i>sodium hypochlorite</i>			
San-I-King No. 451	100 ppm chlorine in solution	-	-

1. Thoroughly clean all surfaces and rinse with potable water prior to treatment. 2. Requires acid activator. 3. After treatment, rinse with potable water.

9. SMALL-SCALE SPRAYER TECHNOLOGY

9.1 Spraying Small Strawberry Plantings

On many small-scale strawberry plantings, spraying often requires special attention to calibration, calculating amounts of pesticide to use, and measuring pesticide products.

To ensure even distribution throughout the canopy, a systematic approach to spraying the whole canopy is essential. Take particular care to cover the top of the canopy as well as ensuring adequate penetration into the inside and middle of the canopy and the fruiting zone. Water sensitive cards (Syngenta) or Surround, kaolin clay, (Engelhard) may be used as tracers to monitor spray distribution.

PRIOR TO SPRAYING—CALIBRATING SPRAYERS

Calibration of backpack sprayers

Use clean water

DYNAMIC CALIBRATION

1. Select correct nozzle and pressure.
2. Measure and mark off an area 10 feet x 10 feet on concrete.
3. Fill sprayer to a known level, mark the fill level.
4. Spray the area on the concrete.
5. Refill sprayer to the fill mark.
6. Compare quantity collected with nozzle chart and desired amount.

STATIC CALIBRATION

1. Select correct nozzle and pressure.
2. Measure and mark off an area 10 feet x 10 feet on concrete.
3. Spray the area and record time taken.
4. Carry out stationary run of same time duration, catching liquid in a graduated measuring jug.
5. Compare quantity collected with nozzle chart and desired amount.

CALCULATING THE AMOUNT OF PESTICIDE TO USE

Some organically approved pesticides are typically sold for large-scale plantings and give application rates on a per acre basis. When converting a known quantity per acre to spray a smaller area, the first step is to measure the area to be sprayed using a tape measure. Divide the number of square feet you have measured by 43,560 to obtain the acreage (in decimal form).

Example:

1. If you are going to spray 20,000 sq. ft,
20,000 divided by 43,560 = 0.459 acre
2. The label states 3 pints of product per acre
Multiply the label rate per acre by the decimal for your area
3 pints multiplied by 0.459 = 1.38 pints
3. Remember there are 16 fl oz in 1 pint.

MEASURING SMALL AMOUNTS OF PESTICIDE

The following tables and examples provide information on converting pesticide rate amounts for smaller areas.

Table 9.1.1. How much powder or granules should I use?

Volume of liquid	100 gallons	25 gallons	5 gallons	1 gallon
Amount of powder or granules to use	4 oz	1 oz	$\frac{3}{16}$ oz	$\frac{1}{2}$ tsp
	8 oz	2 oz	$\frac{3}{8}$ oz	1 tsp
	1 lb	4 oz	$\frac{7}{8}$ oz	2 tsp
	2 lb	8 oz	1 $\frac{3}{4}$ oz	4 tsp
	3 lb	12 oz	2 $\frac{3}{8}$ oz	2 Tbsp
	4 lb	1 lb	3 $\frac{1}{4}$ oz	2 Tbsp + 2 tsp

Powders and granules

Example: The label states 3 lb of powdered product per 100 gallons but you only wish to use a backpack sprayer with a 5-gallon tank. Table 8.1 shows you need to mix in $2\frac{3}{8}$ oz of powder. Use clean weighing scales to provide the correct amount of powder, NEVER use a volumetric measure, e.g. a measuring cup, because the bulk density of different products varies.

Table 9.1.2. How much liquid should I use?

<i>Volume of liquid</i>	100 gallons	25 gallons	5 gallons	1 gallon
Amount of liquid to use	1 gal	2 pts	6 ½ oz	1 ¼ oz
	4 pts	1 pt	3 ¼ oz	⁵ / ₈ oz
	2 pts	½ pt	1 ⁹ / ₁₆ oz	⁵ / ₁₆ oz
	1 ½ pt	6 oz	1 ¼ oz	¼ oz
	1 pt	4 oz	⁷ / ₈ oz	³ / ₁₆ oz
	8 oz	2 oz	⁷ / ₁₆ oz	½ tsp
	4 oz	1 oz	¼ oz	¼ tsp

Table 9.1.3. Dilution of liquid products to various concentrations

<i>Dilution rate</i>	1 gallon	3 gallon	5 gallon
1 in 100	2 Tbsp + 2 tsp	½ cup	³ / ₄ cup + 5 tsp
1 in 200	4 tsp	¼ cup	6 ½ Tbsp
1 in 800	1 tsp	1 Tbsp	1 Tbsp + 2 tsp
1 in 1000	³ / ₄ tsp	2 ½ tsp	1 Tbsp + 1 tsp

Liquids

Example: The label states 4 pts of a liquid product per 100 gallons of spray but you only wish to use a backpack sprayer with a 5-gallon tank. Table 8.2, below, shows you need to mix in 3¼ fl oz of liquid product. Use a clean measuring cylinder or vessel to provide the correct amount of liquid.

Measuring equipment.

Always use measuring equipment that is dedicated only for pesticide use. For very small quantities of liquids, a syringe can be useful. For powder or granular products use weighing scales, do not rely on a measuring cup as the bulk density of products varies.

Safety.

Be sure to wear the proper protective clothing and equipment as required on the pesticide label. Always be aware of watercourses, neighboring properties and changes in the weather.

9.2 Selecting a Small Sprayer for the Small, Organic Strawberry Planting

There are many important points to consider before purchasing a sprayer, not the least of which is the area to spray, the proximity of the local supplier, standard of manufacture, etc. There are many growers with small plantings who need spraying equipment ranging from backpack sprayers to small truck- or ATV-mounted machines.

CANOPY SPRAYERS**Backpack sprayers**

Small capacity (4-5 gallon) sprayers will produce up to approximately 100 psi pressure. Weight is an important consideration and growers should select a sprayer with good, wide, padded straps to ease the load on your shoulders. Correct nozzle selection according to the target is very important to ensure even coverage. A good-sized filling hole at the top is also important.

There are three factors affecting application rate - forward speed, pressure, and nozzle tip size. Unfortunately most inexpensive backpack sprayers have no pressure gauge. Pay more money and purchase a backpack sprayer with a pressure gauge or, better still, purchase a spray management valve as standard or as an option. Normally output increases or decreases according to the pressure in the system, (which is dependent upon how vigorous you are in pumping the handle up and down). A spray management valve, such as a CF valve, will ensure a constant output irrespective of hand pump action. The CF valve evens out fluctuations in pressure, e.g. will only allow a maximum and minimum pressure thus ensuring even flow. The Fountainhead Group sells a backpack sprayer with a simple valve which ensures the correct pressure is not exceeded.

An alternative to the hand-operated backpack sprayer is an electrically-operated backpack sprayer, which utilizes a small rechargeable battery. Maximum pressure is relatively low and it is easier than using a traditional hand pump system, particularly if you have many rows of plants to spray. Similarly a small backpack sprayer fitted with a small gas engine is available. The electric version is quieter to use, but you must remember to recharge the batteries otherwise spraying will be delayed.

Portable mist and air blower backpacks

These are ideal where canopy penetration is required, e.g. denser, vigorous plantings. A small gas engine drives a fan blower which creates an airstream which passes along a hand-held tube (similar to a leaf blower). The tube has a nozzle situated at the end so that liquid spray can be squirted into the airstream. The operator directs the spray cloud towards the canopy by pointing the hand-held tube. It is preferable to point the tube backwards to avoid walking into the spray cloud. Engine speed can be reduced which enables a slower airspeed to match a smaller canopy in early season. They are very good at rustling the canopy and getting good penetration and deposition. They are heavy! Noise is a problem, so ear protection must be worn.

Portable engine-driven gas sprayers

If weight is a problem, and ground conditions are relatively smooth, a number of manufacturers offer a sprayer with a small gas engine and a 10 to 12 gallon tank. Larger capacity tanks (14 to 100 gallons) are often trailed and can be pulled by a lawn tractor, ATV, Gator, or small tractor.

Small, mounted sprayers

Ideal for mounting onto the carrier rack of an ATV, 15 to 25 gallons, they use a small electric pump to provide up to 70 psi. When used with a hand wand and a hose, they can be used to spray short rows. The same system is ideal for weed control and spot spraying of weeds.

Large, skid mounted sprayers

Ideal for fitting into the back of a pick-up truck, these sprayers have a tank capacity of 35 to 200 gallons, and an electric-start gas engine.

HERBICIDE OR GROUND APPLICATION SPRAYERS

Backpack, small ATV-mounted tank, and hand-lance sprayers

These sprayers can be used for herbicide application **BUT** be very careful that there is no carry-over from herbicide residues in the sprayer, therefore wash them out very thoroughly before using them to apply materials other than herbicides. Alternatively, have dedicated herbicide-only equipment.

Controlled Droplet Applicators (CDA)

The use of CDA's will considerably reduce the need to carry vast amounts of water. A spinning disc (battery powered) will produce 95% of the same-size droplets, thus reducing herbicide rates by at least 50% and water rates by 75%. Herbi and Mantis (trade names) are both hand-held CDA sprayers. ATV- or tractor-mounted shielded CDA sprayers such as the Environmist also reduce spray rates while shielding the plants from the spray.

Wick wipers

Where occasional weeds and access over wet land are a problem, the use of a hand-held wick wiper is an easy-to-use, effective option. A small tank, usually contained in the handle, holds the liquid, which soaks a rope wick or a sponge. The rope or sponge can then be wiped against the weeds.

For further information on pesticide application technology visit [Cornell's Pesticide Application Technology](http://Cornell's Pesticide Application Technology website) website.

10. PESTICIDES MENTIONED IN THIS PUBLICATION

Table 10.1 Fungicides, and Bactericides		
Trade Name	Active Ingredient	EPA Reg. No.
Acoidal	sulfur	62562-4
Actinovate-AG	<i>Streptomyces lydicus</i> WYEC 108	73314-1
Agricure	potassium bicarbonate	70870-1
Badge X2	copper oxychloride, copper hydroxide	80289-12
Basic Copper 53	copper sulfate	45002-8
BIO-TAM	<i>Trichoderma asperellum</i> , <i>Trichoderma gamsii</i>	80289-9-69592
BIO-TAM 2.0	<i>Trichoderma asperellum</i> , <i>Trichoderma gamsii</i>	80289-9
Champ WG	copper hydroxide	55146-1
Cinnerate	cinnamon oil	exempt 25(b) pesticide
CS 2005	copper sulfate pentahydrate	66675-3
Cueva Fungicide Concentrate	copper octanoate	67702-2-70051
Defend DF	sulfur	62562-8
Double Nickel 55	<i>Bacillus amyloliquefaciens</i> str. D747	70051-108
Double Nickel LC	<i>Bacillus amyloliquefaciens</i> str. D747	70051-107
Glacial Spray Fluid	mineral oil	34704-849
GrasRoots	cinnamon oil	exempt - 25(b) pesticide
JMS Stylet Oil	paraffinic oil	65564-1
Kaligreen	potassium bicarbonate	11581-2

Table 10.1 Fungicides, and Bactericides		
Trade Name	Active Ingredient	EPA Reg. No.
M-Pede	<i>potassium salts of fatty acids</i>	10163-324
Microthiol Disperss	<i>sulfur</i>	70506-187
Micro Sulf	<i>sulfur</i>	55146-75
Mildew Cure	<i>cottonseed, corn and garlic oils</i>	exempt - 25(b) pesticide
Milstop	<i>potassium bicarbonate</i>	70870-1-68539
Nordox 75 WG	<i>cuprous oxide</i>	48142-4
Nu-Cop 50 WP	<i>copper hydroxide</i>	45002-7
Nu-Cop 50 DF	<i>copper hydroxide</i>	45002-4
Nu-Cop HB	<i>cupric hydroxide</i>	42750-132
Nuke Em	<i>citric acid</i>	exempt 25(b) pesticide
Optiva	<i>Bacillus subtilis</i>	69592-26
Organocide 3-in-1	<i>sesame oil</i>	exempt - 25(b) pesticide
OxiDate 2.0	<i>hydrogen dioxide, peroxyacetic acid</i>	70299-12
PERpose Plus	<i>hydrogen peroxide/dioxide</i>	86729-1
Prestop	<i>Gladiolium catenulatum</i>	64137-11
PureSpray Green	<i>petroleum oil</i>	69526-9
Regalia	<i>Reynoutria sachalinensis</i>	84059-3
RootShield Granules	<i>Trichoderma harzianum Rifai str. T-22</i>	68539-3
RootShield PLUS+ Granules	<i>Trichoderma harzianum str. T-22, Trichoderma virens str. G-41</i>	68539-10
RootShield PLUS+ WP	<i>Trichoderma harzianum str. T-22, Trichoderma virens str. G-41</i>	68539-9
RootShield WP	<i>Trichoderma harzianum Rifai str. KRL-AG2</i>	68539-7
Serenade ASO	<i>Bacillus subtilis str. QST 713</i>	69592-12 and 264-1152
Serenade MAX	<i>Bacillus subtilis str. QST 713</i>	69592-11 and 264-1151
Serenade Opti	<i>Bacillus subtilis str. QST 713</i>	264-1160
Serenade Soil	<i>Bacillus subtilis str. QST 713</i>	69592-12
Serifel	<i>Bacillus amyloliquefaciens</i>	71840-18
Sil-Matrix	<i>potassium silicate</i>	82100-1
Soil Gard	<i>Gladiolium virens str. GL-21</i>	70051-3
SuffOil-X	<i>petroleum oil</i>	48813-1-68539
TerraClean 5.0	<i>hydrogen dioxide, peroxyacetic acid</i>	70299-13
Thiolux	<i>sulfur</i>	34704-1079
Trilogy	<i>neem oil</i>	70051-2
TriTek	<i>mineral oil</i>	48813-1
Zonix Biofungicide	<i>Rhamnolipid Biosurfactant</i>	72431-1

Table 10.2. Insecticides and Miticides

Trade Name	Active Ingredient	EPA Reg. No.
Acoidal	<i>sulfur</i>	62562-4
Aza-Direct	<i>azadirachtin</i>	71908-1-10163
AzaGuard	<i>azadirachtin</i>	70299-17
AzaMax	<i>azadirachtin</i>	71908-1-81268
AzaSol	<i>azadirachtin</i>	81899-4
Azatrol EC	<i>azadirachtin</i>	2217-836
Azera	<i>azadirachtin, pyrethrins</i>	1021-1872
Biobit HP	<i>Bacillus thuringiensis subsp. kurstaki</i>	73049-54
BioLink	<i>garlic juice</i>	exempt 25(b) pesticide

Table 10.2. Insecticides and Miticides		
Trade Name	Active Ingredient	EPA Reg. No.
BioLink Insect & Bird Repellant	<i>garlic juice</i>	exempt 25(b) pesticide
BioRepel	<i>garlic oil</i>	exempt 25(b) pesticide
Cedar Gard	<i>cedar oil</i>	exempt 25(b) pesticide
Cinnerate	<i>cinnamon oil</i>	exempt 25(b) pesticide
Defend DF	<i>sulfur</i>	62562-8
Deliver	<i>Bacillus thuringiensis subsp. kurstaki</i>	70051-69
DES-X	<i>insecticidal soap</i>	67702-22-70051
Dipel DF	<i>Bacillus thuringiensis subsp. kurstaki</i>	73049-39
Ecotec	<i>rosemary and peppermint oil</i>	exempt 25(b) pesticide
Ecozin Plus 1.2% ME	<i>azadirachtin</i>	5481-559
Entrust	<i>spinosad</i>	62719-282
Entrust SC	<i>spinosad</i>	62719-621
Envirepel 20	<i>garlic juice</i>	exempt 25(b) pesticide
Garlic Barrier AG+	<i>garlic juice</i>	exempt 25(b) pesticide
GC-Mite	<i>cottonseed, clove and garlic oils</i>	exempt 25(b) pesticide
Glacial Spray Fluid	<i>mineral oil</i>	34704-849
Grandevo	<i>Chromobacterium subtsugae str. PRAA4-1</i>	84059-17
Javelin WG	<i>Bacillus thuringiensis subsp. kurstaki</i>	70051-66
JMS Stylet Oil	<i>paraffinic oil</i>	65564-1
M-Pede	<i>potassium salts of fatty acids</i>	10163-324
Micro Sulf	<i>sulfur</i>	55146-75
Microthiol Disperss	<i>sulfur</i>	70506-187
Molt-X	<i>azadirachtin</i>	68539-11
Neemix 4.5	<i>azadirachtin</i>	70051-9
Nuke Em	<i>citric acid</i>	exempt 25(b) pesticide
Oleotrol-I	<i>soybean oil</i>	exempt 25(b) pesticide
Omni Supreme Spray	<i>mineral oil</i>	5905-368
Organocide 3-in-1	<i>sesame oil</i>	exempt 25(b) pesticide
PFR-97 20% WDG	<i>Isaria fumosorosea Apopka str. 97</i>	70051-19
PureSpray Green	<i>petroleum oil</i>	69526-9
PyGanic EC 1.4 II	<i>pyrethrins</i>	1021-1771
PyGanic EC 5.0 II	<i>pyrethrins</i>	1021-1772
Safer Brand #567 Pyrethrin & Insecticidal Soap Concentrate II	<i>pyrethrins & potassium salts of fatty acids</i>	59913-9
Sil-Matrix	<i>potassium silicate</i>	82100-1
SuffOil-X	<i>petroleum oil</i>	48813-1-68539
Trilogy	<i>neem oil</i>	70051-2
TriTek	<i>petroleum oil</i>	48813-1
Xentari	<i>(Bacillus thuringiensis subsp. Aizawai, str. ABTS-1857)</i>	73049-40

Table 10.3. Herbicides		
Trade Name	Active Ingredient	EPA Reg. No.
AXXE Broad Spectrum Herbicide	<i>ammonium nonanoate</i>	70299-23

Table 10.4. Mollusk Control Chemicals		
Trade Name	Active Ingredient	EPA Reg. No.
Bug-N-Sluggo	<i>iron phosphate and spinosad</i>	67702-24-70051
Sluggo-AG	<i>iron phosphate</i>	67702-3-54705
Sluggo Slug & Snail Bait	<i>iron phosphate</i>	67702-3-70051

Table 10.5 Sanitizers mentioned in this publication		
Trade Name	Active Ingredient	EPA Reg. No.
CDG Solution 3000	<i>chlorine dioxide</i>	75757-2
Enviroguard Sanitizer	<i>hydrogen peroxide/ peroxyacetic acid</i>	63838-1-527
Oxine	<i>chlorine dioxide</i>	9804-1
Oxonia Active	<i>hydrogen peroxide/ peroxyacetic acid</i>	1677-129
Peraclean 5	<i>hydrogen peroxide/ peroxyacetic acid</i>	54289-3
Peraclean 15	<i>hydrogen peroxide/ peroxyacetic acid</i>	54289-4
Perasan 'A'	<i>hydrogen peroxide/ peroxyacetic acid</i>	63838-1
Per-Ox	<i>hydrogen peroxide/ peroxyacetic acid</i>	833-4
Pro Oxine	<i>chlorine dioxide</i>	9804-9
SaniDate 5.0	<i>hydrogen peroxide/ peroxyacetic acid</i>	70299-19
SaniDate 12.0	<i>hydrogen peroxide/ peroxyacetic acid</i>	70299-18
San-I-King No. 451	<i>sodium hypochlorite</i>	2686-20001
Shield-Brite PAA 5.0	<i>Peroxy acetic acid/ hydrogen peroxide</i>	70299-19-64864
Shield-Brite PAA 12.0	<i>hydrogen peroxide/ peroxyacetic acid</i>	70299-18-64864
StorOx 2.0	<i>hydrogen peroxide/ peroxyacetic acid</i>	70299-7
Tsunami 100	<i>hydrogen peroxide/ peroxyacetic acid</i>	1677-164
Victory	<i>hydrogen peroxide/ peroxyacetic acid</i>	1677-186
VigorOx15 F & V	<i>hydrogen peroxide/ peroxyacetic acid</i>	65402-3
VigorOxLS-15	<i>hydrogen peroxide/ peroxyacetic acid</i>	65402-3

10.1 Pesticides Registered for use in Organic Strawberry Production

At the time the guide was released, the pesticides listed in this guide were allowable for organic production under the National Organic Program (NOP) regulations as set forth in [7 CFR Part 205, sections 600-606](#) and registered for use in New York. The authors relied mainly on the [Organic Materials Review Institute](#) (OMRI) list for pesticides to include. Always check with your certifier before using any new pesticide.

Given the high cost of many pesticides and the limited efficacy data available for many of them, the importance of developing an integrated approach based on cultural practices for disease and insect management, as described in the previous section, cannot be emphasized strongly enough. **Pesticides should not be relied on as a primary method of pest control.** Scouting, forecasting, or trapping pests are important for detecting infestations at an early stage. When conditions do warrant an application, proper choice of materials, proper timing, and excellent spray coverage are essential.

10.2 Pesticide Regulatory Considerations

Organic production focuses on cultural, biological, and mechanical techniques to manage pests on the farm, but in some cases pesticides, which include repellents, allowed for organic production are needed. Pesticides mentioned in this organic production guide are registered by the United States Environmental Protection Agency (EPA) or meet the EPA requirements for a “minimum risk” pesticide. At the time of publication, the pesticides mentioned in this guide also meet New York State Department of Environmental Conservation (NYS DEC) registration requirements for use in New York State. See Cornell’s [Product, Ingredient, and Manufacturer System website](#) for pesticides currently registered for use in NYS. Additional products may be available for use in other states.

To maintain organic certification, products applied must also comply with the National Organic Program (NOP) regulations as set forth in [7 CFR Part 205, sections 600-606](#). The [Organic Materials Review Institute](#) (OMRI) is one organization that reviews products for compliance with the NOP regulations and publishes lists of compliant products, but other entities also make product assessments. Organic growers are not required to use only OMRI listed materials, but the list is a good starting point when searching for allowed pesticides.

Finally, farms grossing more than \$5,000 per year and labeling products as organic must be certified by a NOP accredited certifier who must approve any material applied for pest management. ALWAYS check with the certifier before applying any pest control products. Some certifiers will review products for NOP compliance.

Note that "home remedies" may not be used. Home remedies are products that may have properties that reduce the impact of pests. Examples of home remedies include the use of beer as bait to reduce slug damage in strawberries or dish detergent to reduce aphids on plants. These materials are not regulated as pesticides, are not exempt from registration, and are therefore not legal to use.

Do you need to be a certified pesticide applicator? The Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) defines two categories of pesticides: general-use and restricted-use. NYS DEC also defines additional restricted-use pesticides. Pesticide applicator certification is required to purchase and use restricted-use pesticides. Restricted-use pesticides mentioned in this guide are marked with an asterisk (*). Farmers who purchase and use only general-use pesticides on property they own or rent do not need to be certified pesticide applicators. However, we do encourage anyone who applies pesticides to become certified.

Worker Protection Standard training. If the farm has employees who will be working in fields treated with a pesticide, they must be trained as workers or handlers as required by the federal Worker Protection Standard (WPS). Having a pesticide applicator certification is one of the qualifications needed to be a WPS trainer. Certified pesticide applicators meet the WPS training requirements. For more information on the Worker Protection Standard see: [How To Comply with the Worker Protection Standard](#). See [Revisions To the Worker Protection Standard](#) for a summary of new worker protection standards that will take effect January 2017. Find more information on pesticide applicator certification from the list of [State Pesticide Regulatory Agencies](#) or, in New York State, see the Cornell Pesticide Management Education Program website at <http://psep.cce.cornell.edu>.

10.3 Optimizing Pesticide Effectiveness

Information on the effectiveness of a particular pesticide against a given pest can sometimes be difficult to find. Some university researchers include pesticides approved for organic production in their trials; some manufacturers provide trial results on their web sites; some farmers have conducted trials on their own. Efficacy ratings for pesticides listed in this guide were summarized from university trials and are only provided for some products.

In general, pesticides allowed for organic production may kill a smaller percentage of the pest population, could have a shorter residual, and may be more quickly broken down in the environment than synthetic pesticides. Read the pesticide label carefully to determine if water pH or hardness will negatively impact the pesticide's effectiveness. Use of a surfactant may improve organic pesticide performance. OMRI lists adjuvants in OMRI Products List, Web Edition, [http://www.omri.org/ubersearch/results/adjuvant?type\[\]=opd_listed_product&type\[\]=opd_prohibited_product&type\[\]=opd_removed_product&d\[\]=16997&rb\[\]=17013](http://www.omri.org/ubersearch/results/adjuvant?type[]=opd_listed_product&type[]=opd_prohibited_product&type[]=opd_removed_product&d[]=16997&rb[]=17013).

Regular scouting and accurate pest identification are essential for effective pest management. Thresholds used for conventional production may not be useful for organic systems because of the typically lower percent mortality and shorter residual of pesticides allowed for organic production. When pesticides are needed, it is important to target the most vulnerable stages of the pest. Thoroughly cover plant surfaces, especially in the case of insecticides, since many must be ingested to be effective. The use of pheromone traps or other monitoring or prediction techniques can provide an early warning for pest problems, and help effectively focus scouting efforts.

Pesticide resistance may develop in pathogens, insects, mites, etc. following repeated exposure to the same or similar mode-of-action materials and result in reduced or complete loss of pesticide efficacy against the resistant pest. During the growing season and across growing seasons, pesticides of one mode-of-action should be alternated with those of different modes-of-action to lower the risk of pests developing resistance to the pesticides. See the product label for more information.

11. REFERENCES AND RESOURCES (all links accessed April 2016)

- A & L Eastern Agricultural Laboratories, Inc. 2006. al-labs-eastern.com/
- Agri Analysis, Inc. 2009. www.agrianalysis.com/
- Agricultural Analytical Services Laboratory. Pennsylvania State University. 1998. aasl.psu.edu.
- Agro-One. 2015. <http://dairyone.com/analytical-services/agronomy-services/about-agro-one/>
- Analytical Laboratory and Maine Soil Testing Service, University of Maine. anlab.umesci.maine.edu/

- Berry Diagnostic Tool. Pritts, M. and Heidenreich, C. Cornell University. www.fruit.cornell.edu/berrytool/
- Berry Soil and Nutrient Management - A Guide for Educators and Growers. Pritts, M., Heidenreich, C., McDermott, L. and Miller, J. (eds). 2015. NE SARE 174 pp. <http://www.fruit.cornell.edu/berry/production/soilnutrientmgmt/pdfs/BerrySoilandNutrientManagementGuide.pdf>.
- Buffer pH to Derive Lime Guidelines. 2010. Ketterings, Q., Rao, R., Dietzel, K., and Ristow, P. <http://nmsp.cals.cornell.edu/publications/factsheets/factsheet48.pdf>
- Building Soils for Better Crops, 3rd edition. 2010. Magdoff, F., and VanEs, H. The Sustainable Agriculture Network, Beltsville, MD. 294 pp. <http://www.sare.org/Learning-Center/Books/Building-Soils-for-Better-Crops-3rd-Edition>
- Compendium of Strawberry Diseases, Second Edition. 1998. Maas, J., Editor. American Phytopathological Society. www.apsnet.org/apsstore/shopapspress/Pages/41949.aspx
- Cornell Fruit Resources www.fruit.cornell.edu
- Cornell University Fruit Resources: Berries. www.fruit.cornell.edu/berry/
- Cornell University Insect Diagnostic Laboratory. <http://idl.entomology.cornell.edu/>
- Cornell University Pest Management Guidelines for Berry Crops. Updated yearly. <http://store.cornell.edu/c-875-guidelines.aspx>
- Cornell University Plant Disease Diagnostic Clinic www.plantclinic.cornell.edu/
- Cornell Soil Health Test soilhealth.cals.cornell.edu/
- Cover Crops for Vegetable Growers, Cover Crop Decision Tool. 2010. Björkman, T. Cornell University. <http://covercrops.cals.cornell.edu/decision-tool.php>
- Diseases, Pests, and Beneficial Organisms of Strawberry, Raspberry, and Blueberry. 2013. Lambert, L., Carisse, O., Laplante, G. Hand Vincent, C. <http://www.apsnet.org/apsstore/shopapspress/Pages/02301.aspx>
- Elements of IPM for Strawberries in New York. 2000. NYS IPM Program, Cornell University. <http://www.nysipm.cornell.edu/elements/strawb.asp> EPA Office of Prevention, Pesticides, and Toxic Substances. (2005).
- How To Comply with the Worker Protection Standard for Agricultural Pesticides: What Employers Need to Know. Unit 2: An Introduction to the Worker Protection Standard. 16 pp. <https://www.epa.gov/pesticide-worker-safety/pesticide-worker-protection-standard-how-comply-manual>
- FDA Food Safety Modernization Act (FSMA). www.fda.gov/Food/GuidanceRegulation/FSMA/
- FIFRA Regulation 40 CFR Part 152.25(b) Exemptions for Pesticides of a Character Not Requiring FIFRA Regulation. http://www.ecfr.gov/cgi-bin/text-idx?SID=28b2c435558c77fb3a8074144e32ab3e&mc=true&node=pt40.24.152&rgn=div5#se40.24.152_125
- Good Agricultural Practices (GAPs) Network for Education and Training, Educational Materials. <http://www.gaps.cornell.edu/educationalmaterials.html>
- The Mid Atlantic Berry Guide. 2013-2014. Rudisill, A. Ed. Penn State College of Agricultural Sciences. pubs.cas.psu.edu/freepubs/MAberryGuide
- Minimum Risk Pesticides and Criteria for FIFRA 25(b) Exemption. U.S. Environmental Protection Agency. <http://www2.epa.gov/minimum-risk-pesticides>
- National Organic Program. United States Department of Agriculture: Agricultural Marketing Service (USDA: AMS). <http://www.ams.usda.gov/AMSV1.0/ams.fetchTemplateData.do?template=TemplateA&navID=NationalOrganicProgram&leftNav=NationalOrganicProgram&page=NOPNationalOrganicProgramHome&acct=AMSPW> The National List of Allowed and Prohibited Substances, National Organic Program. Electronic Code of Federal Regulations. www.ecfr.gov/cgi-bin/text-idx?c=ecfr&sid=fbbd316a3eb4c0f243da74a9942b07d8&rgn=div7&view=text&node=7%3A3.1.1.9.32.7.354&idno=7&cc=ecfr
- National Pesticide Information Center: State Pesticide Regulatory Agencies. Cooperative agreement between Oregon State University and the U.S. Environmental Protection Agency. <http://npic.orst.edu/mlrDetail.html?lang=en&to=SPE&state=NY#statePesticide>
- National Sustainable Agriculture Information Service (formerly ATTRA). 2009. National Center for Appropriate Technology (NCAT). attra.ncat.org/organic
- Network for Environment and Weather Applications (NEWA). Cornell University. newa.cornell.edu/
- New York Berry News. Pritts, M., ed. Cornell University. www.fruit.cornell.edu/nybn/
- New York State Berry Growers Association. Cornell University. www.hort.cornell.edu/grower/nybga/
- New York State IPM Fact Sheets for Berries. nysipm.cornell.edu/factsheets/berries/

- New York State Fruit IPM. nysipm.cornell.edu/fruits/
- New York State Pesticide Product, Ingredient, and Manufacturer System (PIMS). 2009. Cooperative Extension, Cornell University. pims.psurl.cornell.edu/
- New York Vineyard Site Evaluation System. 2009. Cornell University and the Institute for the Application of Geospatial Technology. www.nyvineyardsite.org/.
- Northeast Cover Crop Handbook. 1994. Sarrantonio, M. Rodale Institute, PA. Online purchase www.amazon.com/Northeast-Cover-Handbook-Health-Series/dp/0913107174
- Northeast Organic Farming Association of New York. www.nofany.org/
- Nursery Guide for Berry and Small Fruit Crops. 2013. Cornell University. www.fruit.cornell.edu/berry/nurseries/
- OMRI Products List. omri.org/OMRI_products_list
- Organic Farming. National Sustainable Agriculture Information Service (formerly ATTRA). National Center for Appropriate Technology (NCAT). attra.ncat.org/organic
- Organic Farming Development/ Assistance. Sarah Johnston. NYS Dept. of Agriculture and Markets. www.agriculture.ny.gov/AP/organic/
For a list of accredited organic certifiers in NYS – click on “How to Become Certified Organic” and then follow the link “Organizations Providing Organic Certification Services”.
- Organic Materials Review Institute (OMRI). www.omri.org
- Organic Weed Management Website weedecology.css.cornell.edu/manage/
- Penn State Agronomy Guide 2015-2016. Department of Agronomy, The Pennsylvania State University extension.psu.edu/agronomy-guide
- Pesticide Application Technology. Cornell University. web.entomology.cornell.edu/landers/pestapp/
- Pesticide Management Education Program (PMEP). 2008. Cooperative Extension, Cornell University. pmep.cce.cornell.edu/
- Produce Safety Alliance. Dept. of Food Science, Cornell Univ. <http://producesafetyalliance.cornell.edu/alliance> Reducing Deer Damage to Home Gardens and Landscape Plantings, Paul D. Curtis and Milo E. Richmond, Department of Natural Resources, Cornell University, Ithaca, New York 14853. <http://www2.dnr.cornell.edu/ext/info/pubs/Wildlife/reducing%20deer%20damage.pdf>
- Rodale Institute: Leaders in Organic Solutions. www.rodaleinstitute.org/
- Season-Long Strawberry Production with Everbearers for Northeastern Producers EB 401. Lantz, W., Swartz, H., Demchak, K. and Frick, S. http://www.sare.org/content/download/65370/918209/Everbearing_Strawberry_Guide.pdf
- Comprehensive Assessment of Soil Health Website. 2015. Cornell University. soilhealth.cals.cornell.edu/
- Soil Health Testing <http://soilhealth.cals.cornell.edu/> \
- Soil and Plant Tissue Testing Laboratory. 2016. University of Massachusetts. www.umass.edu/soiltest/
- Strawberries: Organic Production. 2007. ATTRA. attra.ncat.org/attra-pub/summaries/summary.php?pub=13
- Strawberry Production Guide for the Northeast, Midwest and Eastern Canada. 1998. NRAES publication #88. Available for purchase from palspublishing.cals.cornell.edu/nra_order.taf?function=detail&pr_booknum=nraes-88
- Title 7. National Organic Program Regulations – from the Electronic Code of Federal Regulations. www.ecfr.gov/cgi-bin/text-idx?c=ecfr&tpl=/ecfrbrowse/Title07/7cfr205_main_02.tpl
- Using Manure and Compost as Nutrient Sources for Fruit and Vegetable Crops. 2005. Rosen, C. J. and Bierman, P. M. University of Minnesota. <http://www.extension.umn.edu/garden/fruit-vegetable/using-manure-and-compost/Using>
- Organic Nutrient Sources. 2009. Sánchez, E., Richard, R. Penn State University. pubs.cas.psu.edu/FreePubs/pdfs/ui256.pdf
- Weeds Of The Northeast. 1997. Uva, R., J. Neal, and J. DiTomaso. Cornell University Press. 397 pp.

11. GLOSSARY

(Adapted from: Wikipedia, www.wikipedia.org/, the free online encyclopedia)

Adjuvant – any substance added to the spray tank, (separate from the pesticide) that will improve the performance of the pesticides, (herbicides, insecticides, miticides, fungicides, bactericides), fertilizers etc. by reducing the surface tension of the water and improving spread and coverage.

Agroecosystem – all of the living and non-living components, including inputs and outputs, that comprise a spatial and functional coherent unit of agricultural activity.

Allelopathy – condition in which one plant emits substances that affect germination, development or growth of other plants in contact with the substance.

Annual – a plant that completes its life cycle within one year (germination, flowering, seed production, death).

Biennial – a flowering plant that takes two years to complete its biological life cycle.

Buffer zone – a physical space of sufficient size that separates two or more areas of activity so that these areas do not affect each other.

Cation exchange capacity – (CEC) is the capacity of a soil to retain and substitute cations (positively charged ions, e.g. potassium) between the soil and the soil solution. CEC is a measure of nutrient retention capacity.

Compost – a combination of plant, animal and other organic materials that have been decomposed largely through aerobic processes into a substance rich in carbon, nutrients, and biological activity.

Crop rotation – the practice of growing, in the same area, in sequential seasons, a series of dissimilar types of crops to avoid the buildup of pathogens and pests that often occurs when one species is continuously cropped.

Frost pocket – an area where still air, cooled by ground-level radiation, travels downhill, replaces warm air, and accumulates to form pockets of very cold air in depressions, valleys, and hollows.

Green manure – a type of cover crop grown for a specific period of time, then incorporated into the soil to add nutrients and organic matter for soil improvement.

Humus – organic matter that is well-decomposed, stable, and contributes to soil tilth and cation exchange.

Immobilization – is when organic matter decomposes and is absorbed by micro-organisms, therefore preventing it being accessible to plants for periods of time. Immobilization is the opposite of mineralization.

Integrated Pest Management (IPM) – a management strategy aimed at insects, mites, plant diseases, weeds, and other pests that uses a variety of planned, complementary tactics including: mechanical devices, physical devices, genetic resistance, biological control, cultural practices, and chemical treatment. It is an ecological approach with a main goal of significantly reducing or eliminating the use of pesticides while at the same time managing pest populations at an acceptable level.

Macroclimate – refers to the regional climate of a broad agricultural area. It can include an area on the scale of tens to hundreds of kilometers.

Mesoclimate – refers to the climate of a particular planting site and is generally restricted to a space of tens or hundreds of meters.

Microclimate – refers to the specific environment in a small restricted space such as a row of plants or corner of a field.

Mineralization – refers to the process where an organic substance is converted to an inorganic substance that can be taken up by the plant.

Nitrogen assimilation – process by which plants expend energy to take up nitrate and ammonium ions and incorporate them into organic molecules required for growth.

Nitrogen budget – accounting that quantifies the nutrients entering the farm (e.g. fertilizers, manure, legumes crops, soil residual nitrogen) and the nutrients leaving the farm (crop harvest, runoff, leaching, and volatilization) for the purpose of balancing inputs and exports.

Nitrogen fixation – the biological process by which nitrogen gas (N_2) in the atmosphere is converted into ammonium compounds that are used by plants.

Organic certification – a certification process for producers of organic food and products that requires strict adherence to production standards for growing, storing, processing, packaging and shipping.

Perched water table – accumulated water above the level of the local water table because impermeable rock or sediment prevents downward movement of water into the local water table.

Perennial – a plant that completes its life cycle (germination, flowering, seed production) over more than one year.

Summer annual – an annual plant that germinates, flowers, produces seed and dies within the same growing season.

Surfactant – (or wetting agent) a soap-like adjuvant added to water or some other liquid to increase wetting properties by reducing the surface tension of the droplets.

Threshold – the density of a pest (insect, mite, plant disease, weed, etc.) at which a control treatment will provide an economic return.

Tilth – a term describing soil that is friable, crumbly, and not compacted which allows rainfall to penetrate and roots to grow without obstruction.

Wind break – (or shelterbelt) is a planting around the edge of a field consisting of one or more rows of trees or shrubs planted in such a manner as to provide shelter from the wind and to protect soil from erosion.

Winter annual – a plant that germinates in the fall or winter, then flowers, produces seed and dies within one year.